


# Exotica and Dark Matter searches

Michele Gallinaro

*LIP Lisbon*


May 4, 2022

- ✓ Introduction
- ✓ Dark matter
- ✓ Exotica searches





# 2012: A new boson discovery




<http://www.elsevier.com/locate/physletb>







# Standard Model theory of everything?

- Discovery of the Higgs boson marks the triumph of the SM
- However, even with the inclusion of the Higgs boson, **SM is an incomplete theory**




# Tests of the SM






# Beyond the Standard Model

The SM answers many of the questions about the structure of matter. But SM is not complete; still many unanswered questions:

- 
- a) Why do we observe matter and almost no antimatter if we believe there is a symmetry between the two in the universe?
  - b) What is this "dark matter" that we can't see that has visible gravitational effects in the cosmos?
  - c) Are quarks and leptons actually fundamental, or made up of even more fundamental particles?
  - d) Why are there three generations of quarks and leptons? What is the explanation for the observed pattern for particle masses?
  - e) How does gravity fit into all of this?

# Dark matter and energy


- What is that accounts for 96% of the Universe?  
Nobody knows.
- It is one of the greatest mysteries of Science





# What can we look for?

A crowded field. At the LHC we can search for some of these



# How?

- Search for new phenomena
- Look for New Physics
- **Indirect searches**
  - precision measurements, event properties, etc.
- **Direct searches**
  - resonances, specific final states, model-(in)dependent searches, etc.
- Production and decay rates, event characteristics, advanced tools





# Dark Matter

## What is it?

- DM does not interact electromagnetically
- DM interacts gravitationally

Visual map



# Dark Matter (cont.)

Why is it interesting?

- We do not see it...but we feel it

Mass map






# Dark Matter (cont.)


## How do we find DM?

- Need to understand how it interacts with Universe
- Traditionally through a mediator
- Yields at least two new particles





# Searching for DM



## Direct Detection



## Particle Colliders




## Indirect Detection







# How do we find it: @underground

- Through a nuclear recoil




# How do we find it: @Space

- Through annihilation
  - Cosmic rays from DM



# How do we find it: @LHC


- Produced it through a mediator





# DM at the LHC


- CMS/ATLAS experiments **not** designed for DM searches



# DM searches at LHC

How do we find DM at the LHC?

- DM production gives MET signature




# DM searches: backgrounds

What are the backgrounds?

- $Z \rightarrow \nu\nu$

–very similar to signal






# DM searches: backgrounds (cont.)

How to discriminate signal against the background?

- Look for high MET:



Study hadronic recoil

$$MET = -\sum_{All\ particles} p_T$$


$$MET(Z \rightarrow \nu\nu) = - \text{Z recoil} + p_T(\nu\nu)$$

$$MET(Z \rightarrow \nu\nu) = - \text{Z } p_T$$

# DM searches: backgrounds (cont.)


How to discriminate signal against the background?

- Can fit the shape and look for signal




# DM+Z


signal



background




- Main background is from ZZ di-bosons
- Understanding ZZ di-boson pT is critical





# Build a V-tagger

- Two jets are more collimated at high  $p_T$




- At **low  $p_T$**  jets are “resolved”
  - Focus on reconstructing di-jets with mass near  $W$  mass
- At **high  $p_T$**  get one “fat” jet
  - Focus on identifying one jet with mass near  $W$  mass
- Use additional variables to improve discrimination



# DM+jet/V

arXiv:1712.02345

## DM search in mono-jet/V





Need good control of systematics




# VBF: H(invisible)

arXiv:1809.06682, arXiv:1809.05937


- In the SM,  $B(H \rightarrow \text{invisible})$  only 0.1%
  - Any significant deviation would indicate BSM
  - Signature: Large MET,  $\Delta\phi(jj)$ , veto  $\ell/b$ -jets
    - C&C and shape fit of  $m(jj)$
  - Main bkg:  $V + \text{jets}$  (95%)
  - Tag with forward jets+MET
  - Cross section  $\sim 4\text{pb}$
  - Small background
- 



Shape: bkg-only fit in CRs+SR

35.9 fb<sup>-1</sup> (13 TeV)

Set limits:  $B(H \rightarrow \text{inv.}) < 0.37 \text{ (0.28) @95\%CL}$







# DM+Higgs

arXiv:1703.05236

- DM search with  $H(\rightarrow bb, \gamma\gamma)$
- Model dependent search
- $Z'$  2HDM Model







- No significant excess
- Set limits for coupling  $g=0.8$



# Dark Matter



arXiv:1903.01400, arXiv:1901.0155, CMS-EXO-18-009

- Complementarity to direct/indirect searches
- DM particles:
  - interact via spin-0 & spin-1 mediators
  - are undetected (MET) recoiling against SM particle(s)
- Extensive program of mono-X searches (X=jet,  $\gamma$ , lepton, W, Z, t, tt, bb, H)
- No excess observed
- Interpretation through simplified models (DM and mediator masses, couplings to DM and SM)



# Experimental results

- Limits for given couplings between SM and DM interaction
- **Competitive limits at low masses wrt other experiments**





⇒ Collider results complement direct searches for low masses (<5-10GeV)



# Resonant searches




Among the highest dijet mass event recorded:  $m_{jj} = 8.12 \text{ TeV}$

## Resonant Searches



# BSM models predict new resonances



- BSMs predict resonances with spin 0,1,2
- Are quarks fundamental particles?
  - Excited quarks in models of compositeness
- Randall-Sundrum (RS) models
  - Spin-2 graviton (KK-particle)
- Heavy-Vector Triplets
  - Spin-1 resonance
  - Models based on strength of vector boson interactions
- Sequential SM
  - $Z'$  and  $Z$  with same couplings to fermions
  - Width proportional to the mass



# New phenomena in di-jet events


CMS-EXO-17-026, arXiv:1910.08447

- Searches up to high masses
- QCD predicts a smooth, monotonic decrease in dijet invariant mass
- Search for a localized excess
- No significant excess observed






# Searching for dilepton resonances



CMS Experiment at the LHC, CERN

Data recorded: 2017-Jun-27 15:39:36.789504 GMT

Run / Event / LS: 297599 / 134277310 / 86






**Dimuon candidate event:  
Reconstructed mass of 2.4 TeV**



# Di-muon events



- Di-muon events: a re-discovery of the SM



# High-mass dilepton resonances


arXiv:1803.06292, arXiv:1903.06248, CMS-EXO-19-019

- Search for dilepton ( $ee, \mu\mu$ ) resonance




# Search for diboson resonances

- Heavy BSM resonances ( $>1\text{TeV}$ ) may decay into SM bosons (W,Z, H)
- Several final states
- Experimental challenges
  - SM bosons decay mostly to quarks
  - Due to large Lorentz boost, decay products merge into single jet
  - Clustered within a large-cone jet ( $R=0.8$ )
- Look into jet substructure
  - Jet “grooming”: get rid of soft jet components from UE/pileup, keep constituents from hard scatter
  - Apply filters (mass drop, pruning, trimming)



# Diboson resonances

- Many potential final states are possible
  - $WW/WZ$ ,  $ZW/ZZ$ ,  $VV$
- Hadronic channels with high sensitivity in high mass region






# Jet grooming


arXiv:0802.2470

## Mass drop/filtering


- Identify approx. symmetric sub-jets (with smaller mass than sum)



**Mass drop:** define 2 sub-jets



**Filtering:** re-cluster  $j_1, j_2$  constituents




# Jet grooming (cont.)

arXiv:0912.1342, arXiv:0912.0033


## “Trimming”

- Uses kT algorithm to make subjets (subjets with  $p_T^i/p_T < \text{cut}$  removed)



## “Pruning”

- Recombine jet constituents, while veto wide-angle/softer constituents



# W, Z, H reconstruction



arXiv:1707.01303

- Grooming and jet mass

- Pruning
- soft drop (stable w/pileup, and good jet mass resolution  $\sim 10\%$ )

- Vector boson tagging ( $V \rightarrow qq$ )


- n-subjettiness  $\tau_{21}$ : how consistent with 2 sub-jets
- Categorization according to purity: high ( $<0.35$ ) and high ( $>0.35$ )



# W, Z, H reconstruction (cont.)

arXiv:1707.01303

- Higgs boson tagging ( $H \rightarrow b\bar{b}$ )
  - Double b-tagging
  - Exploit b-tagging to identify two b-quarks in same jet
  - Soft-lepton information
  - Combines tracking and vertexing in MVA







# Searching for diboson resonances

arXiv:1906.08589




- No significant excess in any of the observed final states
- Exclusion limits: HVT models excluded up to 4.1 TeV, Spin-2 RS models up to 2.8 TeV
- Large improvements due to new methods for jet reconstructions and boson tagging



# Search for multi-lepton final states

JHEP 03(2020)051

- Search for new heavy particles
  - Heavy fermions/scalars produced in association with  $t\bar{t}$
- Search for 3 or more lepton final states
- Pair production of  $W/Z/H \rightarrow \Sigma\Sigma$ 
  - Scalar sum of lepton  $p_T$  ( $L_T$ )
  - Bin and count ( $L_T + \text{MET}$ )



# $X \rightarrow VV \rightarrow qqqq$

arXiv:1708:05379



- All hadronic resonance search with single (qV) or double (VV) V-tag
  - At least 2 back-to-back jets  $p_T > 200 \text{ GeV}$
  - Categorization (jet mass,  $\tau_{21}$ )
- Background estimation: “bump hunt” fit data with power law



Candidate ZZ event  
Dijet mass: 3.2 TeV

Anti- $k_T$  R=0.8 jet  
 $p_T$  1321 GeV  
 $\eta$  -0.40  
 $\phi$  -2.71  
 $M_{SD}$  103 GeV  
 $\tau_{21}$  0.23



Anti- $k_T$  R=0.8 jet  
 $p_T$  1374 GeV  
 $\eta$  0.79  
 $\phi$  0.43  
 $M_{SD}$  94.8  
 $\tau_{21}$  0.29



# $X \rightarrow VH \rightarrow qqbb$

arXiv:1707.01303

- All-hadronic search for  $V \rightarrow qq$  and  $H \rightarrow bb$  resonances
  - dedicated identification for  $H \rightarrow bb$  (b-tagging)
- Use categories
  - V-jet mass (W or Z), V-jet  $\tau_{21}$  (high-purity, low-purity), H-jet (tight and loose b-tag)






- Similar topology and background estimate to VV resonance search
- No significant excess found in data



# $X \rightarrow ZV \rightarrow \ell\ell qq$

arXiv:1803.10093

- Search for resonances in  $Z \rightarrow ee/\mu\mu$ ,  $V \rightarrow qq$
- Clean final state (leptons)
  - Good mass resolution, good efficiency
- $\tau_{21}$  categorization (HP, LP)
- Parametrize main bkg ( $Z$ +jets), fit to data in sidebands, take shape from MC





- Data compatible with SM-only hypothesis

# $X \rightarrow WV \rightarrow \ell \nu qq$

arXiv:1802.09407, B2G-19-002

- Search for a resonance decaying to  $WV$  in lepton+jet channel
- Categorization in  $\tau_{21}$  and  $W/Z$  mass
- Sideband+transfer function for bkg estimate





- Similar sensitivity to  $Z(\ell)V(qq)$  search
- Excluded up to 1.1-3.1 TeV

# $X \rightarrow VH \rightarrow \ell \nu qq$



PLB 768(2017)137

- Search for a resonance decaying to VH in leptonic channels
  - $Z \rightarrow \nu\nu$ : transverse mass  $m_T(VH)$
  - $W \rightarrow \ell\nu$ : top control region
  - $Z \rightarrow \ell\ell$ : high-efficiency dilepton ID
  - $H(bb)$  b-tagging
- Sideband bkg prediction

- Heavy vector triplet ( $Z'$ ,  $W'$ )
- $g_V, g_H$  ( $c_V, c_F$ ): couplings



# Combination of diboson searches







# Combination of resonance searches

PLB 768(2019)134952

- Combination of searches for heavy resonances decaying to boson and lepton final states
- Large gain in statistical combination






# Resonance searches: Summary



# Leptoquarks

ATLAS-CONF-2021-008



- Pair production of leptoquarks
  - Study  $LQ \rightarrow b\tau$  final state
  - Can be interpreted in SUSY
- Main background from top quark pair events in different final states, determined from CRs
- Signal: use  $M_{T2}$  (stransverse mass) and  $s_T$  (scalar sum of  $\tau$  and jet  $p_T$ ) variables



# Heavy lepton resonances

EPJC 81(2021)218, arXiv:2202.02039

- Two resonances in one process:  $W \rightarrow NL$
- Off-shell  $W$  decays to two new particles
- Signal selection:
  - OS and SS lepton pairs possible
  - two jets and MET
- Backgrounds from simulations and data CRs







# Vector-like quarks

## Motivation

- Simplest extension allowed in the quark sector
- Spin  $\frac{1}{2}$  fermions with vector coupling
- Can mix with SM quarks and modify their couplings to the  $W/Z/H$  bosons
- Sizeable mixing with 3<sup>rd</sup> family,  $b$  and  $t$

## Properties



- Produced via strong and EWK interactions
- Mainly pair-produced
- Both CC and NC decay modes



# VLQ searches

PLB 779(2018)82



- Search for VLQ pair production decaying to  $WbWb$
- Search in the **boosted regime**
- Can reconstruct the VLQ system



# Looking forward: PPS


CERN-LHC-2014-021

- The Precision Proton Spectrometer is a joint CMS and TOTEM project that aims at measuring the surviving **scattered protons** on both sides of CMS in standard running conditions
- **Tracking** and **timing** detectors inside the beam pipe at ~210m from IP5
- Project approved in Dec. 2014 by LHCC
- Data taking started in 2016 (full scope from 2017)



# PPS physics motivations

- **Central Exclusive Production**
  - photon-photon collisions
  - gluon-gluon fusion in color singlet,  $J^{PC}=0^+$
- **High- $p_T$  system in central detector, together with very forward protons in PPS**
  - momentum balance between central system and forward protons, provides strong kinematical constraints
  - Mass of central system measured by momentum loss of the two leading protons
- **Gauge boson production by photon-photon fusion and anomalous couplings** ( $\gamma\gamma WW$ ,  $\gamma\gamma ZZ$ , and  $\gamma\gamma\gamma\gamma$ )
- **Search for new BSM resonances**
- **Study of QCD in a new domain**






# Detectors



- Tracking detectors

- Goal: measure proton momentum
- Technology: silicon 3D pixels (6 planes per pot)




- Timing detectors

- Goal: identify primary vertex, reject “pileup”
- $\sigma_{\text{time}} \sim 10\text{ps} \Rightarrow \sigma_z \sim 2\text{mm}$
- Technology: silicon/diamond



# Roman Pot insertion

- Insertion procedure validated in 2016 by the LHC
  - Improvements carried out wrt earlier versions (RF shielding, cylindrical pots, ferrite, copper coating)
- Minimum distance of approach dramatically affects detector acceptance and physics reach
- A few mm ( $\sim 15\sigma$ ) from beam in nominal high-luminosity runs
  - Monitor beam losses, showers, interplay with collimators, beam impedance (heating, vacuum and beam orbit stability)





# LHC tunnel @ PPS location

215m

CT-PPS  
timing


214m

CT-PPS  
tracking

beam

220m

CT-PPS  
tracking







# Timing detectors

Use timing to reject pileup background

- Two scenarios studied:
  - 10ps and 30ps time resolution
- Baseline: solid state detectors
- Detector options investigated:
  - Diamond sensors
  - Fast silicon sensors (UFSD, HFS)
- Status:
  - Diamond and LGAD detectors installed







# WW production

JHEP 08(2016)119


- Study of process:  $pp \rightarrow pWWp$ 
  - Clean process: W in central detector and “nothing” else, intact protons can be detected far away from IP
  - Exclusive production of W pairs via photon exchange: QED process, cross section well known
- Backgrounds:
  - inclusive WW,  $\tau\tau$ , exclusive two-photon  $\gamma\gamma \rightarrow ll$ , etc.
- Events:
  - WW pair in central detector, leading protons in PPS
- SM observation of WW events
- Anomalous coupling study
  - AQGCs predicted in BSM theories
  - parameters:  $a_0^W/\Lambda^2$ ,  $a_c^W/\Lambda^2$
- Deviations from SM can be large





# Exclusive Dileptons

JHEP 07(2018)153

- Study exclusive processes at the EWK scale
- Search for two-photon production of opposite charge lepton pair with forward proton tagging



signal





Background: SD, DD, DY, dibosons, PU

- Signal selected with:
- at least one proton tagged, muons, kinematic selection

# Exclusive Dileptons (cont.)

JHEP 07(2018)153

- Correlation between the  $\xi$  values in central system vs RP
- $12\mu\mu$ ,  $8ee$  candidates observed ( $>5\sigma$  over expected bkg)
- First observation of two-photon production of a lepton pair at this mass range




# AQGC expected limits

arXiv:1607.03745

Expected limits @95%CL:

$$\begin{aligned} a_0^W / \Lambda^2 &= 2 \times 10^{-6} \quad (3 \times 10^{-6}) \\ a_C^W / \Lambda^2 &= 7 \times 10^{-6} \quad (10 \times 10^{-6}) \end{aligned}$$









# $\gamma\gamma \rightarrow \gamma\gamma$ : Anomalous couplings, etc.


PRD 89(2014)114004, CMS-EXO-18-014

- Indirect search: neutral quartic gauge couplings (forbidden in SM) in  $\gamma\gamma \rightarrow \gamma\gamma$
- Expect to provide best sensitivity at LHC
- Sensitive to axion-like particles



# Physics w/ forward protons


JHEP 07(2018)153, CMS-PRO-21-001, CMS-TOP-21-007, CMS-SMP-21-004



## Proton reconstruction

CMS-PRO-21-001


PPS collected more than 100/fb of data in Run2



## Exclusive top quark pairs

CMS-TOP-21-007


Search for central exclusive production of  $t\bar{t}$  pairs in pp interactions with tagged protons



## $Z\gamma + X$ production

CMS-PAS-EXO-19-009


Search for anomalous  $Z/\gamma^*$  central production with 2017 data



## Exclusive $WW/ZZ$



CMS-SMP-21-004

Search for  $\gamma\gamma \rightarrow WW/ZZ$  with forward protons




# Prospects for Run3 and beyond

- More luminosity in a more challenging environment
- Will enhance the mass reach in the search for new particles
- Need to meet experimental challenges
  - Aging of detector, improve/adapt capability
  - Integrated luminosity: 300-3000/fb
  - peak luminosity of  $2 \times 10^{35} \text{cm}^{-2}\text{s}^{-1}$
  - pileup will be  $\sim 150$  or higher (Phase2)
  - large radiation doses





# BSM searches: resonances, etc.






# HL-LHC upgrades

Luminosity of  $\sim 3000 \text{ fb}^{-1}$  expected for HL-LHC

- Tracking information in “L1 track-trigger”
  - Tracker designed to enable finding all tracks w/ $p_T > 2 \text{ GeV}$  in  $< 4 \mu\text{s}$
- Tracker is all silicon but with much higher granularity, up to  $|\eta|=4$ 
  - $> 2$  billion pixels and strips
- High Granularity Endcap Calorimeters
  - Sampling of EM showers: every  $\sim 1\lambda$  (28 samples) w/pixels, and every  $\sim 0.35\lambda$  (24 samples) with pixels+scintillator to map 3D shower development
  - $\sim 6 \text{ M}$  channels in all
- Precision timing to add a 4<sup>th</sup> dimension to object reconstruction



# Future: HL-LHC upgrades

## Trigger/HLT/DAQ

- Track information in hardware event selection
- 750 kHz hardware event selection
- 7.5 kHz events registered

## Barrel EM calorimeter

- New electronics
- Low operating temperature  $\approx -10^\circ$

## Muon systems

- New DT & CSC electronics
- New chambers  $1.6 < \eta < 2.4$
- Muon tagging  $2.4 < \eta < 3$


## New Endcap Calorimeters

- Rad. Tolerant
- 5D measurement




## New Tracker

- Rad. Tolerant - light
- High Definition measurement
- 40 MHz selective readout for hardware trigger
- Extended Pixel coverage to  $\eta \approx 3.8$


Beam radiation and luminosity  
Common systems and infrastructure



# Precision Timing Layer



- Time-of-flight precision  $\sim 30\text{ps}$ 
  - $|\eta| < 3$ ,  $p_T > 0.7\text{GeV}$
  - Crystal+SiPM: rad hard to  $2 \times 10^{14} n_{eq} \text{cm}^{-1}$
- Provide  $\sim x4-5$  effective PU reduction
  - 15% merged vertices reduce to 1.5%
  - Low PU track purity of vertices recovered
- Showers timed to 30ps in calorimeters





# Summary

- Excellent consistency of SM but **SM is incomplete**
- Direct and indirect searches for New Physics
  - Collected  $\sim 150/\text{fb}$  @13 TeV in 2015-2018
  - $\sim 300/\text{fb}$  to be collected in the next few years (up to LS3)
- Many studies performed with data collected so far
  - New dedicated algorithms being developed
  - Dark Matter, Exotica, signature-based searches
  - Other BSM searches
- Searches provide **no hints for BSM yet**

