





Tutorial on

Data Analysis



FCT Fundação para a Ciência e a Tecnologia
MINISTÉRIO DA CIÊNCIA, TECNOLOGIA E ENSINO SUPERIOR




LABORATÓRIO DE INSTRUMENTAÇÃO
E FÍSICA EXPERIMENTAL DE PARTÍCULAS

goals

perform a simple data analysis

- visualise the data
- manipulate data ntuples
- produce, process, and display data histograms
 - select different physics signals
 - plot kinematic distributions, inspect detector/trigger effects
- extract physics parameters from data
 - measure signal yields by performing a likelihood fit
 - inspect statistical and systematic errors
 - (extra) perform a differential measurement

Detector & Event Reconstruction & Visualisation



calorimeters:

measure particle's energy by absorbing it

trackers:

detect trajectory of charged particles

muons:

detected in outer detector layers

CMS

3.8T Superconducting Solenoid

Lead tungstate
E/M Calorimeter (ECAL)

Hermetic ($|\eta|<5.2$)
Hadron Calorimeter (HCAL)
[scintillators & brass]

All Silicon Tracker
(Pixels and Microstrips)


Redundant Muon System
(RPCs, Drift Tubes,
Cathode Strip Chambers)

CMS

3.8T Superconducting Solenoid

Hermetic ($| $n|<5.2$)
Hadron Calorimeter (HCAL)
[scintillators & brass]$

Lead tungstate
EM Calorimeter (ECAL)




All Silicon Tracker
(Pixels and Microstrips)



Redundant Muon System
(RPCs, Drift Tubes,
Cathode Strip Chambers)

Key:


- Muon
- Electron
- Charged Hadron (e.g. Pion)
- - - Neutral Hadron (e.g. Neutron)
- - - Photon



a di-electron event



Event Display of a Candidate Electron-Positron Pair with an Invariant Mass of 2.9 TeV





di-photons

Experiment at the LHC, CERN


Data recorded: 2015-Nov-02 21:34:00.662277 GMT,

Run / Event / LS: 260627 / 854678036 / 477



$m_{\gamma\gamma} \sim 750 \text{ GeV}$



[CMS-PHO-EVENTS-2015-007](#)




di-jets




$X \rightarrow jj$



a di-muon event




$$X \rightarrow \mu\bar{\mu}$$




the di-muon analysis



the di-muon spectrum ($X \rightarrow \mu\mu$)

50 years of particle physics in one plot!






from detector to physics ...



di-muon ‘invariant mass’ ?



particle identification


- signal in muon chambers
→ it's a muon!
- ⇒ $m = m(\mu) \sim 106 \text{ MeV}/c^2$

particle trajectory

- muon chambers but especially the silicon tracker
- ⇒ linear momentum, $\mathbf{p} \equiv (p_x, p_y, p_z)$


- ⇒ form 4-momentum of each muon: $\mathbf{P}_\mu \equiv (E, p_x, p_y, p_z)$
- ⇒ that of the di-muon pair $\mathbf{P}_{\mu\mu} = \mathbf{P}_{\mu 1} + \mathbf{P}_{\mu 2} = \mathbf{P}_{\text{x} \rightarrow \mu\mu}$
- ⇒ invariant mass $\mathbf{P}_{\mu\mu} \cdot \mathbf{P}_{\mu\mu} = \mathbf{M}_{\mu\mu}^2 = (\mathbf{M}_{\text{x}})^2$

the reconstructed di-muon spectrum




feature: variable bin widths, resolution-dependent, properly normalized, doubly-log scales

fit the data



- inspect quality of fit
 - can model be improved?
 - hint: final state radiation ($\mu \rightarrow \mu\gamma$) may distort shape
- establish a **fit model**
 - signal; Gaussian
 - background: polynomial
- extract **signal parameters**
 - yield ($N \pm \sigma_N$), mass ($m \pm \sigma_m$)
- estimate **systematic errors**
 - does the choice of fit model affect the measured results ?
 - quantify the systematic variations by employing different models
- quote **final measurements**
 - $N \pm \sigma_{\text{stat}} \pm \sigma_{\text{syst}}$


what's the physics process ?



production: strong force

decay: electroweak force

what are the peaks?



Check their measured properties from: <http://pdglive.lbl.gov>

Z $J = 1$

See related reviews:

[Z Boson](#)

PDF

[Anomalous ZZ \$\gamma\$, Z \$\gamma\gamma\$, and ZZV Couplings](#)

PDF

[Anomalous W/Z Quartic Couplings \(QGCs\)](#)

PDF

1

55. Z Boson**55. Z Boson**

Revised August 2018 by M. Grünwald (University Coll. Dublin) and A. Gurtu (CERN; TIFR Mumbai).

Precision measurements at the Z -boson resonance using electron–positron colliding beams began in 1989 at the SLC and at LEP. During 1989–95, the four LEP experiments (ALEPH, DELPHI, L3, OPAL) made high-statistics studies of the production and decay properties of the Z . Although the SLD experiment at the SLC collected much lower statistics, it was able to match the precision of LEP experiments in determining the effective electroweak mixing angle $\sin^2\theta_W$ and the rates of Z decay to b - and c -quarks, owing to availability of polarized electron beams, small beam size, and stable beam spot.

The Z -boson properties reported in this section may broadly be categorized as:

- The standard ‘lineshape’ parameters of the Z consisting of its mass, M_Z , its total width, Γ_Z , and its partial decay widths, $\Gamma(\text{hadrons})$, and $\Gamma(\ell\bar{\ell})$ where $\ell = e, \mu, \tau, \nu$;
- Z asymmetries in leptonic decays and extraction of Z couplings to charged and neutral leptons;
- The b - and c -quark-related partial widths and charge asymmetries which require special techniques;
- Determination of Z decay modes and the search for modes that violate known conservation laws;
- Average particle multiplicities in hadronic Z decay;
- Z anomalous couplings.

The effective vector and axial-vector coupling constants describing the Z -to-fermion coupling are also measured in $p\bar{p}$ and ep collisions at the Tevatron and at HERA. The corresponding cross-section formulae are given in Section 39 (Cross-section formulae for specific processes) and Section 16 (Structure Functions) in this *Review*. In this minireview, we concentrate on the measurements in e^+e^- collisions at LEP and SLC.

The standard ‘lineshape’ parameters of the Z are determined from an analysis of the production cross sections of these final states in e^+e^- collisions. The $Z \rightarrow \nu\bar{\nu}(\gamma)$ state is identified directly by detecting single photon production and indirectly by subtracting the visible partial widths from the total width. Inclusion in this analysis of the forward-backward asymmetry of charged leptons, $A_{FB}^{(0,\ell)}$, of the τ polarization, $P(\tau)$, and its forward-backward asymmetry, $P(\tau)^{fb}$, enables the separate determination of the effective vector (\bar{g}_V) and axial vector (\bar{g}_A) couplings of the Z to these leptons and the ratio (\bar{g}_V/\bar{g}_A), which is related to the effective electroweak mixing angle

C $I(J^P) = 0(1/2^+)$ Charge = $\frac{2}{3} e$ Charm = +1

▶ Expand all sections

Z MASS

 $91.1876 \pm 0.0021 \text{ GeV}$ $1.27 \pm 0.02 \text{ GeV}$

c-QUARK MASS

 $11.76^{+0.05}_{-0.10}$ m_c/m_s MASS RATIO 4.58 ± 0.01 m_b/m_c MASS RATIO $3.45 \pm 0.05 \text{ GeV}$ $m_b - m_c$ QUARK MASS DIFFERENCE

INSPIRE search

 $J/\psi(1S)$ $I^G(J^{PC}) = 0^-(1^{--})$ J/ $\psi(1S)$ MASS $3096.900 \pm 0.006 \text{ MeV}$ J/ $\psi(1S)$ WIDTH $92.6 \pm 1.7 \text{ keV (S = 1.1)}$ J/ $\psi(1S)$ Decay Modes

▶ Expand all decays

Mode	Fraction (Γ_i / Γ)	Scale Factor/ Conf. Level	P(MeV/c)
Γ_1 hadrons	(87.7 ± 0.5)%		
Γ_2 virtual $\gamma \rightarrow$ hadrons	(13.50 ± 0.30)%		
Γ_3 ggg	(64.1 ± 1.0)%		
Γ_4 γgg	(8.8 ± 1.1)%		
Γ_5 e^+e^-	(5.971 ± 0.032)%	1548	
Γ_6 $e^+e^-\gamma$	[1] (8.8 ± 1.4) × 10 ⁻³	1548	
Γ_7 $\mu^+\mu^-$	(5.961 ± 0.033)%	1545	

▶ Decays involving hadronic resonances

▶ Decays into stable hadrons

▶ Radiative decays

▶ Dalitz decays

▶ Weak decays




▶ Charge conjugation (C), Parity (P), Lepton Family number (LF) violating modes

▶ Other decays


Cross section

“N=L.σ”

$$\frac{d^2\sigma(Q\bar{Q})}{dp_T dy} \mathcal{B}(Q\bar{Q} \rightarrow \mu^+ \mu^-) = \frac{N_{fit}(Q\bar{Q})}{\mathcal{L} \cdot \mathcal{A} \cdot \epsilon \cdot \Delta p_T \cdot \Delta y}$$



an effective area of interaction
unit: barn, $1\text{b} = 10^{-28} \text{ m}^2 = 100\text{fm}^2$



- N: fitted signal yield
- A: detector acceptance from simulation
- E: detector reconstruction and trigger efficiencies (simulation or data-driven)
- L: integrated sample luminosity

LHC Open Data

opendata
CERN

- the LHC collaborations make good chunks of their data publicly available
 - <http://opendata.cern.ch/>
- along with tools & software & examples
- for data visualisation and analysis
- from event reconstruction algorithms to machine learning challenges
- via virtual machines (with no need to install different software packages)
- few pointers
 - <http://opendata.cern.ch/visualise/events/cms>
 - <http://www.i2u2.org/elab/cms/event-display/>
- you're invited to **explore the LHC data** also on your own leisure

