


SWG0 present and (near) future



Mário Pimenta
September 2025




> 90 institutes, 16 countries

Argentina, Brazil, Chile, **China**, Croatia, Czech Republic, France, Germany, Italy, Mexico, Netherlands, Peru, **Portugal**, South Korea, United Kingdom, United States

SWGO, 2024: the year of decisions

Baseline Layout and Detectors




$\sim 1 \text{ Km}^2$

Core

+

Outer Array



Mexico City
April 2024

Double Layer
metallic WCD

Mercedes
rotomolded WCD

Site

Pampa La Bola, AAP, Chile
4,770 m a.s.l.




Rio,
July 2024



SWGO near Future

Path Finder

2025/26





First activity @ site

2+1 metal D tanks,
2+1 plastic Mercedes tanks
(one with isolation and diffusive walls)





SWGO-A



core array $\sim 20\,000\text{ m}^2$

USA (NSF – 17M US\$) +
Germany, ...


SWGO-H (or PEV array...)



outer array 1 – 4% FF

Brazil, Italy +
Portugal, Czech Republic, Chile,...

SWGO Science vision




Meeting agenda

SWGO-LIP meeting

📅 Friday Sep 19, 2025, 2:00 PM → 5:20 PM Europe/Lisbon

📍 LIP-Lisboa/3-311 - Sala de Seminários (LIP Lisboa)

- | | | | |
|--|-----------|--|-------|
| 2:00 PM | → 2:10 PM | Introduction and agenda | 🕒 10m |
| Speaker: Prof. Mário Pimenta (LIP) | | | |
| 2:10 PM | → 2:30 PM | Mercedes Rotomolded WCD | 🕒 20m |
| Speaker: Prof. Ulisses Barres de Almeida (CBPF) | | | |
| 2:30 PM | → 2:50 PM | WCD thermal simulations | 🕒 20m |
| Speaker: Prof. Luis Filipe Mendes (IST) | | | |
| 2:50 PM | → 3:10 PM | A proposal for a low-energy trigger | 🕒 20m |
| Speaker: Pedro Costa (LIP/IST) | | | |
| 3:10 PM | → 3:30 PM | Progress in outdoor sealed RPCs | 🕒 20m |
| Speaker: Alberto Blanco (LIP) | | | |
| 3:30 PM | → 3:50 PM | Coffee break | |
| 3:50 PM | → 4:10 PM | Machine learning applications for SWGO | 🕒 20m |
| Speaker: Prof. Ruben Conceição (IST/LIP) | | | |
|  Ruben_SWGO-LIP_... | | | |
| 4:10 PM | → 4:30 PM | LCm and Ptail gamma/hadron discriminators | 🕒 20m |
| Speaker: Lucio Gibilisco (LIP) | | | |
| 4:30 PM | → 5:00 PM | General Discussion | |

@ Pampa La Bola !



LIP Roadmap

LIP@Lower energies

- trigger
- pattern recognition
- tagging muon stations
- Alerts? Multi-Messengers?
- ...

LIP@Detector,

- Mercedes (with CBPF)
- Layout optimization
- Thermal simulations
- RPCs@high altitude
- Calibrations?
- IACTs?
- ...


LIP@Higher energies

- LCm
- Ptail
- FR_{\min} , X_{\max} , $X\mu_{\max}$
- Composition
- ...

LIP@ ...

- neutrinos
- hadronic interactions
- outreach
- Pevatrons @ the Galaxy Centre?
- ...

The sensitivity curve



A new variable: LCm

Gamma/hadron discrimination at high energies through the azimuthal fluctuations of air shower particle distributions at the ground

R. Conceição, L. Gíbilisco, M. Pimenta and B. Tomé

Laboratório de Instrumentação e Física Experimental de Partículas (LIP),
Av. Prof. Gama Pinto, 2, 1649-003 Lisbon, Portugal
Departamento de Física, Instituto Superior Técnico (IST), Universidade de Lisboa,
Av. Rovisco Pais, 1, 1049-001, Lisbon, Portugal

E-mail: ruben@lip.pt, gibilisco@lip.pt, pimenta@lip.pt, bernardo@lip.pt

Received April 27, 2022

Revised September 28, 2022

Accepted October 11, 2022

Published October 26, 2022



Abstract. Wide field-of-view gamma-ray observatories must fight the overwhelming cosmic ray background to identify very-high-energy astrophysical gamma-ray events. This work introduces a novel gamma/hadron discriminating variable, LCm , which quantifies the azimuthal non-uniformity of the particle distributions at the ground. This non-uniformity, due to the presence of hadronic sub-showers, is higher in proton-induced showers than in gamma showers. The discrimination power of this new variable is then discussed, as a function of the air shower array fill factor, in the energy range 10 TeV to 1 PeV, and compared to the classical gamma/hadron discriminator based on the measurement of the number of muons at the ground. The results obtained are extremely encouraging, paving the way for the use of the proposed quantity in present and future large ground-array gamma-ray observatories.

Keywords: gamma ray experiments, ultra high energy photons and neutrinos, cosmic ray experiments




ArXiv ePrint: [2204.12337](https://arxiv.org/abs/2204.12337)

$$C_k = \frac{2}{n_k(n_k - 1)} \frac{1}{\langle S_k \rangle} \sum_{i=1}^{n_k-1} \sum_{j=i+1}^{n_k} (S_{ik} - S_{jk})^2$$

$$LCm \equiv \log(C_k)|_{r_k=r_m}$$



It works!







A new variable: P_{tail}^{α}

$$P_{\text{tail},i} = C_{r_i}(S_i) \rightarrow P_{\text{tail}}^{\alpha} = \sum_i^n (P_{\text{tail},i})^{\alpha}$$

Station variable

Event variable



I. INTRODUCTION

The selection, with good efficiency and high purity, of highly energetic gamma rays or the determination of the nature of the charged cosmic rays is one of the major challenges for cosmic ray and gamma ray experiments.

The direct detection of neutral and charged cosmic rays by high-altitude balloons or satellites is enabled at high energies (above tens of TeV for gamma rays and tens of eV for TeV for charged cosmic rays) due to the scarcity of such particles and the limited detection area of such detectors (typically a few m^2) [1]. Thus, the only viable option is indirect detection, achieved by measuring the longitudinal development of the Extensive Air Shower (EAS) produced by the interaction of these particles in the Earth's atmosphere [2, 3], or by studying the distribution of the EAS particles that reach the ground [4, 5].

Several different experimental methods and discriminant variables have been developed to select gamma-ray events from the huge hadronic background and to discriminate between showers that might have been produced by different atomic nuclei (typically from hadrons to iron) [6, 7]. No unique or perfect solution exists, although, above a few TeV, the direct measurement of the number of muons arriving at the ground is widely accepted as the best possible discriminant variable and has indeed allowed the detection of gamma rays with energies up to the PeV by the LHAASO collaboration [8, 9]. More recently, a new gamma/hadron discriminating variable, LC_{m} , based on the measurement of the azimuthal non-uniformity of the particle distributions at the ground



in Water Cherenkov Detectors (WCD) arrays, was introduced [10] and, through simulations, it has been claimed that it might reach equivalent background rejection factors of about 10^4 at energies about 1 PeV [11]. The latter quantity has, however, shown limited discrimination power for composition and hadronic interactions studies.

In this article, we introduce a novel variable denoted as P_{tail}^{α} , designed for WCD ground arrays. By focusing on events falling within a specific energy range at the ground, we construct distributions of signals across stations, categorized into discrete distance bins from the shower core. These distributions can be either derived from available data or generated through simulation when data is lacking. P_{tail}^{α} provides a quantitative measure on an event-by-event basis, indicating the number of stations exhibiting signals within the upper tail of these signal distributions. The rationale behind this variable, inspired by a method developed by the IceTop/IceCube collaboration [12], lies in the observation that in events with comparable reconstructed energy, the signal recorded by the WCD stations tends to be higher when struck by energetic sub-showers. These sub-showers, composed of muons and highly energetic electromagnetic particles, serve as a distinct signature of hadronically-induced showers [13].

The manuscript is organized as follows: In Section II, all the simulation sets are described; In Section III, the new variable, P_{tail}^{α} , is introduced; In Section IV, the correlation of this new variable with the number of muons that hit the WCD stations in gamma, proton or iron events with reconstructed energy between 10 TeV and 1.4 PeV is analyzed; In Section V, the efficiency of the new variable to select high-purity gamma event samples, as well as to determine the nature of charged cosmic rays events is re-

* ghilardi@lip.pt

It works!



Access the number of muons without burying the WCDs !