



Contribution ID: 54

Type: **Workshop 2025/2026**

## Critical Gravitational Collapse with a Charged Scalar Field

*Wednesday 28 January 2026 16:15 (15 minutes)*

In recent groundbreaking work, it was shown that there is a regime of critical collapse where the critical solution is an extremal Reissner-Nordström black hole. It was also conjectured that such critical phenomena would occur for the spherically symmetric Einstein-Maxwell-charged scalar field system. In this work, we report on early developments to test this conjecture. We perform the 3+1 decomposition for the Maxwell equations as well as the massless Klein-Gordon equation. We implement the evolution of the vector potential  ${}^{(3)}A_i$  in the existing numerical relativity code BAMPS, which introduces the curl constraint  $\mathcal{G}_A^i = (D \times A)^i - B^i$ . To check for the correctness of the code, we simulate an electromagnetic dipole in flat background and Reissner-Nordström black hole in fixed background. The first simulation was successful, showing exponential convergence. However, the latter one showed exponential growth of  $\mathcal{G}_A^i$  with resolution and in time. This suggests that constraint damping for  $\mathcal{G}_A^i$  needs to be implemented in the future.

### Field of Research/Work

Cosmology, Astrophysics, and Gravitation

**Author:** YAO, Joana (Instituto Superior Técnico)