




Detectando Partículas no LHC


Nuno Leonardo
nuno@cern.ch




a cadeia de aceleradores do CERN, e o LHC



o magneto mais potente




Milhares de milhões de colisões (por segundo)




25 ns bunch crossing
25 ns entre les paquets

the LHC detectors


ATLAS




CMS




LHCb



ALICE






**Solenóide
supercondutor**

**Calorímetro hadrónico
(HCAL)**


**Calorímetro
electromagnético
(ECAL)**

**Traçador
(detectores de silício)**

Camêras de muões




Identificando as partículas estáveis




Derived from CMS Detector Slice from CERN

Identificando as partículas estáveis




Derived from CMS Detector Slice from CERN

Identificando as partículas estáveis



Derived from CMS Detector Slice from CERN

Identificando as partículas estáveis




Charged Hadron signature: Signals in the Tracker and HCAL; almost nothing in ECAL and nothing in the muon chambers.

Charged hadrons, such as protons and pi plus or pi minus (made of pairs of quarks), are bent by the magnetic field and travel straight through the ECAL leaving almost no signals. Upon reaching the HCAL they are slowed to a stop by the dense materials, producing showers of secondary particles along the way that in turn produce light in thin layers of plastic **scintillator** material. The amount of light is proportional to the energy of the incoming hadron.


Derived from CMS Detector Slice from CERN

mas as colisões do LHC não são simples ...


[leading order]




- [higher order corrections] objects are finally reconstructed using information from different detector subsystems combined in a **particle flow** algorithm
 - electrons radiate via bremsstrahlung
 - photons may convert to e^+e^- pairs in the tracker
 - **jet** (q,g) energy is formed of charged/neutral hadrons (65%/10%) and photons (25%): calorimeter and tracker info exploited
 - **missing E_T** requires 'full event' reconstruction



muão + anti-muão + (fundo)




o espectro de massa $\mu^+\mu^-$




A re-descoberta do Modelo Padrão, 2010 (primeiros dados do LHC)


electrão + positrão + (fundo)




o espectro de massa e^+e^-





Dois fótons + (fundo)






?


$m_{??} \sim 700 - 800 \text{ GeV}$



$$\rho = \frac{p}{ZeB}$$





muão + energia-em-falta + (fundo)




- mas ... e os Neutrinos?
- os neutrinos quase que não interagem com a matéria
- São detectados indirectamente, garantindo o balanço de energia (no plano perpendicular ao feixe)
- Os neutrinos correspondem assim à energia em falta ('MET')

CMS Experiment at LHC, CERN
Data recorded: Wed Jul 8 23:50:40 2015 CEST
Run/Event: 251251 / 39089589
Lumi section: 64










MEDIDAS DE PRECISÃO



partículas novas e raras



<http://opendata.cern.ch/visualise/events/CMS>

opendata CERN

ABOUT SEARCH EDUCATION RESEARCH

Home > Education > Visualise Events > CMS


Explore CMS open data and visualise detector events

Need HELP?

iSpy WebGL

Detector

- Pixel Barrel
- Pixel Endcap (+)
- Pixel Endcap (-)
- Tracker Inner Barrel
- Tracker Outer Barrel
- Tracker Inner Detector (+)
- Tracker Inner Detector (-)
- Tracker Endcap (+)
- Tracker Endcap (-)



Education


Visualise events, check reconstructed data, run tools or build your own!

Start learning



Research

Get the genuine working environments, virtual machines and datasets to start your research


Start analysing



iSpy-Online




- obtendo a distribuição de massa dos candidatos Z





EvNo	E1	px1	py1	pz1	pt1	eta1	phi1	Q1	E2	px2	py2	pz2	pt2	eta2	phi2	Q2	M
128943239	72.89895	13.36098	-26.087	66.74727	29.3095	1.5612	-1.09746	1	37.6277	-10.9181	35.80517	-3.82334	37.3966	-0.10197	1.86677	-1	90.31227

- Busca de acontecimentos raros
- e.g. $H \rightarrow ZZ$ com $Z \rightarrow e^+ e^-$ e $Z \rightarrow \mu^+ \mu^-$



- Acontecimentos $H \rightarrow \gamma\gamma$
 - Um dos canais de descoberta do Higgs



Depósitos de energia no ECAL,
Sem deixar sinal do traçador
(detectores de silício)

