

Search for new interactions in the top quark sector

1st IGFAE/LIP Workshop | Braga · May 4, 2018



Maura Barros · Nuno Castro · Ana Peixoto



Universidade do Minho
Escola de Ciências

FCT Fundação para a Ciéncia e a Tecnologia
MINISTÉRIO DA CIÉNCIA, TECNOLOGIA E ENSINO SUPERIOR · Portugal

COMPETE 2020 **PORTUGAL 2020**




Motivation

FCNC limits overview



1

- ▶ Flavour Changing Neutral Currents (FCNC): change in the fermion flavour through the emission of a neutral boson
- ▶ Many types of New Physics (NP) lead to FCNC, often at tree level



- ▶ More than 10 orders of magnitude left for $\text{FCNC } \mathcal{BR}(t \rightarrow u|cZ)$ before Standard Model (SM) suppression

Signal Simulation



- ▶ MC production via **left-handed utZ** and **ctZ** tensor couplings (K_{ut}^L , K_{ct}^L) at LO QCD using TopFCNC UFO model within **MADGRAPH 5**
- ▶ EFT Lagrangian using **dimension-6 operators** in LO QCD

$$\mathcal{L}_{\text{LO}} = \mathcal{L}_{\text{SM}} + \sum_{q \in \{u,c\}} \left[\begin{array}{l} \frac{g_s}{2m_t} \bar{q} T^a \sigma^{\mu\nu} (\zeta_{qt}^L P_L + \zeta_{qt}^R P_R) t G_{\mu\nu}^a + \frac{e}{2m_t} \bar{q} \sigma^{\mu\nu} (\lambda_{qt}^L P_L + \lambda_{qt}^R P_R) t A_{\mu\nu} \\ + \frac{g_w}{2c_w} \bar{q} \gamma^\mu (X_{qt}^L P_L + X_{qt}^R P_R) t Z_\mu + \frac{g_w}{4c_w M_Z} \bar{q} \sigma^{\mu\nu} (K_{qt}^L P_L + K_{qt}^R P_R) t Z_{\mu\nu} \\ + \frac{1}{\sqrt{2}} \bar{q} (\eta_{qt}^L P_L + \eta_{qt}^R P_R) t H \end{array} \right] + h.c.$$

Motivation


$q t Z$ anomalous coupling




3

FCNC $u|c+g \rightarrow tZ$:

- ▶ In decay: $t\bar{t}$ production
- ▶ In production: single-top
- ▶ Clean 3ℓ signature: $e^+e^-|\mu^+\mu^- + e|\mu + b\text{-jet} + E_t$
- ▶ utZ and ctZ anomalous couplings considered
- ▶ Main backgrounds: $t\bar{t}$, $t\bar{t}Z$, Z +jets and diboson processes



tZ production via FCNC



$t\bar{t}$ decay via FCNC



Analysis with decay mode

Summary

4



- ▶ $t\bar{t}$ production where a top quark decays to a **W boson** and a **b-quark** and the other decays to a **Z boson** and an **up or charm quark**
- ▶ Final state with **three charged leptons** and at least **two jets** where one of them is *b*-tagged
- ▶ One signal region and **five control** regions focusing on the $t\bar{t}Z$, WZ , ZZ processes



<https://arxiv.org/abs/1803.09923>


Analysis with production mode

Choice of discriminant variable



5

- ▶ tZ production particularly **sensitive** to utZ coupling
- ▶ **Harder distribution** of the **transverse momentum** of the Z boson compared with the background processes



- ▶ For a low value of $m_T(\ell, \nu)$, the $Z+jets$ background can be **isolated**
- ▶ **Two complementary** regions separated by **low** and **high** $m_T(\ell, \nu)$ regions defined

Signal and Control Regions

Event selection



6

- ▶ Three **signal** regions
- ▶ Two **control** regions: $t\bar{t}$ and WZ diboson backgrounds

Selection	SR			CR	
	m_T^W -low	m_T^W -high	2-jet	$t\bar{t}$ 2	WZ
No. leptons	= 3	= 3	= 3	= 3	= 3
No. jets	= 1	= 1	= 2	≥ 2	= 1
No. b -jets	= 1	= 1	= 1	≥ 1	= 0
Z candidate	≥ 1	≥ 1	≥ 1	< 1	≥ 1
m_T^W	< 50 GeV	> 50 GeV	-	-	-

- ▶ Definition of the **transverse mass** of the W boson:





$$m_T^W = \sqrt{2E_t p_t^\ell (1 - \cos(\Delta\varphi(E_t, p_t^\ell)))}$$

Signal Regions

Post-fit plots



7




Results


95% expected upper limits on cross-sections, couplings and BRs



Limit for K_{ut}^L



Limit for K_{ct}^L



Coupling	$\sigma_{\text{FCNC IZ}} [\text{pb}]$	$K/\Lambda_{\text{NP}} [\text{TeV}^{-1}]$	$\mathcal{BR}(t \rightarrow qZ)$	
			ATLAS	CMS
utZ anomalous coupling				
Expected	0.284	0.0171	2.25×10^{-4}	1.50×10^{-4}
Observed	0.318	0.0183	2.52×10^{-4}	2.40×10^{-4}
ctZ anomalous coupling				
Expected	0.369	0.0547	2.34×10^{-3}	3.70×10^{-4}
Observed	0.401	0.0581	2.54×10^{-3}	4.50×10^{-4}



Limits on $\mathcal{B}(t \rightarrow uZ)$ comparable to CMS results

Interference Studies

Production and Decay



- ▶ FCNC at production and decay can have the **same final state**
- ▶ **Interferences effects** between both modes should be considered when **combined limits** are obtained
- ▶ Feynman diagrams for the decay (left) and production (right)





Interference Studies

Production and Decay



- ▶ Phenomenological study being currently done using Monte Carlo generation with MadGraph5 importing the TopFCNC UFO model
- ▶ Only qtZ anomalous coupling is considered at this stage




Photoproduction via FCNC

Prospects



- ▶ Probing uty and cty couplings
- ▶ Final state composed by proton remnants, a top quark and a proton
- ▶ An interesting possibility is using the data collected by the **ATLAS Forward Proton (AFP)** detector in the forward region



Summary



- ▶ Distinct analyses for decay and production **implemented**

Expected upper limits				
\sqrt{s} [TeV]	Luminosity [fb^{-1}]	Mode	$\mathcal{BR}(t \rightarrow qZ)$	
13	35.9 - CMS	Decay and production	$1.5 \cdot 10^{-4}$ ($q = u$)	
13	35.9 - CMS	Decay and production	$3.7 \cdot 10^{-4}$ ($q = c$)	
13	36.1 - ATLAS	Decay	$2.4 \cdot 10^{-4}$ ($q = u$)	
13	36.1 - ATLAS	Decay	$3.2 \cdot 10^{-4}$ ($q = c$)	
13	36.1 - This analysis	Production	$2.2 \cdot 10^{-4}$ ($q = u$)	
13	36.1 - This analysis	Production	$2.3 \cdot 10^{-3}$ ($q = c$)	

- ▶ Study of the **interferences** between both channels is under way
- ▶ **Photoproduction via FCNC** processes is an interesting possibility for a **future analysis**

Appendix

Motivation

FCNC suppression in the SM



- ▶ Flavour Changing Neutral Currents (FCNC): change in the fermion flavour through the emission of a neutral boson
- ▶ Many types of **New Physics (NP)** lead to FCNC, often at **tree level**
- ▶ Predictions for the branching ratios of **FCNC top decays** by the SM and some NP models

Process	SM	QS	2HDM	FC 2HDM	MSSM	R SUSY
$t \rightarrow g u$	$3.7 \cdot 10^{-14}$	$1.5 \cdot 10^{-7}$	—	—	$8 \cdot 10^{-5}$	$2 \cdot 10^{-4}$
$t \rightarrow Z u$	$8 \cdot 10^{-17}$	$1.1 \cdot 10^{-4}$	—	—	$2 \cdot 10^{-6}$	$3 \cdot 10^{-5}$
$t \rightarrow \gamma u$	$3.7 \cdot 10^{-16}$	$7.5 \cdot 10^{-9}$	—	—	$2 \cdot 10^{-6}$	$1 \cdot 10^{-6}$
$t \rightarrow H u$	$2 \cdot 10^{-17}$	$4.1 \cdot 10^{-5}$	$5.5 \cdot 10^{-6}$	—	$\approx 10^{-5}$	$\approx 10^{-6}$
$t \rightarrow g c$	$4.6 \cdot 10^{-12}$	$1.5 \cdot 10^{-7}$	$\approx 10^{-4}$	$\approx 10^{-8}$	$8 \cdot 10^{-5}$	$2 \cdot 10^{-4}$
$t \rightarrow Z c$	$1 \cdot 10^{-14}$	$1.1 \cdot 10^{-4}$	$\approx 10^{-7}$	$\approx 10^{-10}$	$2 \cdot 10^{-6}$	$3 \cdot 10^{-5}$
$t \rightarrow \gamma c$	$4.6 \cdot 10^{-14}$	$7.5 \cdot 10^{-9}$	$\approx 10^{-6}$	$\approx 10^{-9}$	$2 \cdot 10^{-6}$	$1 \cdot 10^{-6}$
$t \rightarrow H c$	$3 \cdot 10^{-15}$	$4.1 \cdot 10^{-5}$	$1.5 \cdot 10^{-3}$	$\approx 10^{-5}$	$\approx 10^{-5}$	$\approx 10^{-6}$

Common Selection

Event selection



14

- ▶ Common cuts for the objects:

- ▶ $p_t(\ell_1) > 27 \text{ GeV}$
- ▶ $p_t(\ell_2) > 25 \text{ GeV}$
- ▶ $p_t(\ell_3) > 15 \text{ GeV}$
- ▶ $p_t(j) > 30 \text{ GeV}$
- ▶ 1 b -jet (77% WP, $|\eta| < 2.5$)
- ▶ no other jet (77% WP, $|\eta| < 4.5$)
- ▶ $|m(\ell\ell) - m(Z)| < 10 \text{ GeV}$

- ▶ Definition of the transverse mass of the W boson:

$$m_T^W = \sqrt{2E_t p_t^\ell (1 - \cos(\Delta\varphi(E_t, p_t^\ell)))}$$

Signal and Control Regions

Event selection



- ▶ Three **signal** regions
- ▶ Two **control** regions: $t\bar{t}$ and WZ diboson backgrounds

Selection	SR			VR		CR		
	m_T^W -low	m_T^W -high	2-jet	Z+jets 1	Z+jets 2	$t\bar{t}$ 1	$t\bar{t}$ 2	WZ
No. leptons	= 3	= 3	= 3	= 2	= 2	= 2	= 3	= 3
No. jets	= 1	= 1	= 2	= 1	= 2	= 2	≥ 2	= 1
No. b -jets	= 1	= 1	= 1	= 1	= 2	= 2	≥ 1	= 0
Z candidate	≥ 1	≥ 1	≥ 1	≥ 1	≥ 1	< 1	< 1	≥ 1
m_T^W	< 50 GeV	> 50 GeV	-	-	-	-	-	-

Signal Regions

Event yields



16

SR1 m_T^W -low					SR2 m_T^W -high				
	Pre-fit	Post-fit	Ratio		Pre-fit	Post-fit	Ratio		
$tZ K_L^{ut}$	11.5 ± 1.2	11.5 ± 1.2	1.00 ± 0.10	$tZ K_L^{ut}$	20.5 ± 1.5	20.5 ± 1.3	1.00 ± 0.07		
$tZ K_L^{ct}$	12.7 ± 1.3	12.7 ± 1.3	1.00 ± 0.10	$tZ K_L^{ct}$	25.2 ± 2.1	24.9 ± 1.9	0.99 ± 0.07		
tZj	1.5 ± 0.5	1.5 ± 0.5	1.00 ± 0.32	tZj	3.3 ± 1.1	3.4 ± 1.0	1.04 ± 0.31		
$t\bar{t}V, t\bar{t}H$	1.3 ± 0.4	1.3 ± 0.4	1.04 ± 0.32	$t\bar{t}V, t\bar{t}H$	3.2 ± 1.1	3.4 ± 1.0	1.04 ± 0.30		
$t\bar{t}, tW$	8 ± 8	13.5 ± 3.2	1.74 ± 0.41	$t\bar{t}, tW$	13 ± 13	25 ± 5	1.85 ± 0.41		
$Z + \text{jets}$	35 ± 38	52 ± 14	1.46 ± 0.40	$Z + \text{jets}$	0.9 ± 1.0	1.0 ± 0.7	1.20 ± 0.75		
VV	23 ± 7	26 ± 5	1.12 ± 0.20	VV	39 ± 13	46 ± 7	1.19 ± 0.19		
Total	69 ± 40	94 ± 15	1.36 ± 0.22	Total	60 ± 19	78 ± 8	1.32 ± 0.13		
Data	78			Data	83				

SR3 2-jets				
	Pre-fit	Post-fit	Ratio	
$tZ K_L^{ut}$	28.8 ± 2.0	28.4 ± 1.8	0.99 ± 0.06	
$tZ K_L^{ct}$	34.3 ± 2.1	34.3 ± 2.0	1.00 ± 0.06	
tZj	30 ± 9	31 ± 9	1.04 ± 0.30	
$t\bar{t}V, t\bar{t}H$	18 ± 6	19 ± 5	1.04 ± 0.29	
$t\bar{t}, tW$	15 ± 15	27 ± 6	1.80 ± 0.39	
$Z + \text{jets}$	19 ± 20	30 ± 11	1.59 ± 0.57	
VV	48 ± 35	59 ± 17	1.23 ± 0.36	
Total	130 ± 45	166 ± 15	1.28 ± 0.12	
Data	170			

Signal Extraction Fit

Strategy



Use a binned likelihood function with a signal strength, $\mu = \sigma_{\text{FCNC}}/1 \text{ pb}$ (not multiplied by $\mathcal{B}(tZ \rightarrow 3\ell\nu b)$), and nuisance parameters (NPs) for systematic uncertainties. The $p_t(Z)$ distributions in the 3 signal and 2 control regions with 5 bins each used as discriminant variable

- ▶ Find μ such that p-value on B-only Asimov dataset is 0.05
 - ▶ expected upper limit
- ▶ Fit B-only model to B-only Asimov dataset
 - ▶ expected estimates of NPs
- ▶ Fit S+B model to S+B Asimov dataset
 - ▶ expected approximate impact of uncertainties
- ▶ Fit B-only model to parts of observed data
 - ▶ Estimates of NPs for post-fit plots, modelling checks

Signal Extraction Fit


Nuisance Parameters



18

B, Asimov

JET_BJES_Response
 JET_EtaInterpolation_Modeling
 JET_EtaInterpolation_NonClosure
 JET_EtaInterpolation_TotalStat
 JET_Flavor_Composition
 JET_JER_SINGLE_NP
 JET_PtRel_OffsetAll
 JET_PtRel_OffsetTauVBF
 JET_Pileup_PtTerm
 JET_Pileup_RhoTopology
 JET_PunchThrough_MC15
 Lumi
 Pileup_SF
 bTagSF_B_0
 bTagSF_B_1
 bTagSF_B_2
 bTagSF_B_3
 bTagSF_B_4
 bTagSF_B_5
 bTagSF_C_0
 bTagSF_C_1
 bTagSF_C_2
 signal_2l_mm
 signal_2l_mm_generator_rate
 signal_2l_mm_generator_shape
 signal_MtWl_mm
 signal_MtWl_mm_generator_rate
 signal_MtWl_mm_generator_shape
 signal_MtWl_mm_generator_shape
 tt_fake_lepton_pt
 ttbar3_lvv_norm
 ttbar3_lvv_generator_rate
 ttbar3_lvv_generator_shape
 ttv_norm
 tz_norm
 vv_3l_Njet_1
 vv_3l_Njet_2
 vv_3l_Njet_3
 vv_3l_Njet_more
 vv_Njet_BC
 wtb_norm
 wz_vv_norm
 wz_vv_generator_rate
 wz_vv_generator_shape
 zjets_3l_Njet_1
 zjets_3l_Njet_2
 zjets_3l_Njet_3
 zjets_3l_Njet_more
 zjets_Njet_BC
 zjets_fake_lepton_pt
 zjets_fakes_norm
 zjets_norm



B, observed

