

Global Analysis of SSAs from Current and Future Data



Daniel Pitonyak

Lebanon Valley College, Annville, PA, USA



22nd Particles and Nuclei International Conference (PANIC 2021)

September 5, 2021




Background

Leading Twist TMDs



		Quark Polarization		
		Un-Polarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \bullet$		$h_1^\perp = \bullet - \bullet$ Boer-Mulders
	L		$g_{1L} = \bullet \rightarrow - \bullet \rightarrow$ Helicity	$h_{1L}^\perp = \bullet \rightarrow - \bullet \rightarrow$
	T	$f_{1T}^\perp = \bullet \uparrow - \bullet \downarrow$ Sivers	$g_{1T}^\perp = \bullet \uparrow - \bullet \uparrow$	$h_1 = \bullet \uparrow - \bullet \uparrow$ Transversity $h_{1T}^\perp = \bullet \uparrow - \bullet \uparrow$


Leading Twist TMDs




		Quark Polarization		
		Un-Polarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \bullet$		$h_1^\perp = \bullet - \bullet$ Boer-Mulders
	L		$g_{1L} = \bullet \rightarrow - \bullet \rightarrow$ Helicity	$h_{1L}^\perp = \bullet \rightarrow - \bullet \rightarrow$
	T	$f_{1T}^\perp = \bullet \uparrow - \bullet \downarrow$ Sivers	$g_{1T}^\perp = \bullet \uparrow - \bullet \uparrow$	$h_1 = \bullet \uparrow - \bullet \uparrow$ Transversity $h_{1T}^\perp = \bullet \uparrow - \bullet \uparrow$

Naïve time-reversal odd (T-odd)

$$\ell p^\uparrow \rightarrow \ell h X$$




$$\{\pi, p\} p^\uparrow \rightarrow \{\ell^+ \ell^-, W^\pm, Z\} X$$



$$F_{UT}^{\sin(\phi_h - \phi_S)} = \mathcal{C} \left[-\frac{\hat{h} \cdot \vec{k}_T}{M} \mathbf{f}_{1T}^\perp D_1 \right]$$

$$F_{TU}^{\sin \phi} = \mathcal{C} \left[-\frac{\hat{h} \cdot \vec{k}_{aT}}{M_a} \mathbf{f}_{1T}^\perp \bar{f}_1 \right]$$


$$\tilde{f}_{1T}^{\perp(1)}(x, b_T; Q^2, \mu_Q) \sim F_{FT}(x, x; \mu_{b_*}) \exp \left[-S_{pert}(b_*(b_T); \mu_{b_*}, Q, \mu_Q) - S_{NP}^{f_{1T}^\perp}(b_T, Q) \right]$$



OPE
(NLO available from
Scimemi, Tarasov,
Vladimirov (2019))

$g_{f_{1T}^\perp}(x, b_T) + g_K(b_T) \ln(Q/Q_0)$
(Aybat, et al. (2012); Echevarria, et al. (2014))

Leading Twist TMDs



Quark Polarization

		Un-Polarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \text{circle with dot}$		$h_1^\perp = \text{circle with dot} - \text{circle with dot}$ Boer-Mulders
	L		$g_{1L} = \text{two circles with arrows pointing right}$ Helicity	$h_{1L}^\perp = \text{two circles with arrows pointing right}$
	T	$f_{1T}^\perp = \text{circle with up arrow} - \text{circle with down arrow}$ Sivers	$g_{1T}^\perp = \text{two circles with arrows pointing up}$	$h_{1T}^\perp = \text{two circles with arrows pointing up}$ Transversity

Survive
integration
over k_T

Leading Twist TMDs



		Quark Polarization		
		Un-Polarized (U)	Longitudinally Polarized (L)	Transversely Polarized (T)
Nucleon Polarization	U	$f_1 = \bullet$		$h_1^\perp = \bullet - \bullet$ Boer-Mulders
	L		$g_{1L} = \bullet \rightarrow - \bullet \rightarrow$ Helicity	$h_{1L}^\perp = \bullet \rightarrow - \bullet \rightarrow$
	T	$f_{1T}^\perp = \bullet \uparrow - \bullet \downarrow$ Sivers	$g_{1T}^\perp = \bullet \uparrow - \bullet \uparrow$	$h_1 = \bullet \uparrow - \bullet \uparrow$ Transversity $h_{1T}^\perp = \bullet \uparrow - \bullet \uparrow$

Chiral odd


Survive
integration
over k_T

$$\delta q \equiv \int_0^1 dx [h_1^q(x) - h_1^{\bar{q}}(x)] \quad g_T \equiv \delta u - \delta d$$

The tensor charge of the nucleon is one of its fundamental charges and is important for BSM studies in, e.g., beta decays (Gonzalez-Alonso, et al. (2018),...)



Processes sensitive to TMDs can play an important role in these efforts (Courtois, et al. (2015); Yamanaka, et al. (2017), Liu, et al. (2018),...)

Lattice QCD has also calculated the tensor charges with great precision (Gupta, et al. (2018); Hasan, et al. (2019), Alexandrou, et. (2019),...)



$$\ell p^\uparrow \rightarrow \ell h X$$

$$e^+ e^- \rightarrow h_1 h_2 X$$




Collins function

$$F_{UT}^{\sin(\phi_h + \phi_S)} = \mathcal{C} \left[-\frac{\hat{h} \cdot \vec{p}_\perp}{M_h} h_1 \textcolor{blue}{H}_1^\perp \right] \quad F_{UU}^{\cos(2\phi_0)} = \mathcal{C} \left[\frac{2\hat{h} \cdot \vec{p}_{a\perp} \hat{h} \cdot \vec{p}_{b\perp} - \vec{p}_{a\perp} \cdot \vec{p}_{b\perp}}{M_a M_b} \textcolor{blue}{H}_1^\perp \bar{H}_1^\perp \right]$$


$$\tilde{h}_1(x, b_T; Q^2, \mu_Q) \sim h_1(x; \mu_{b_*}) \exp[-S_{pert}(b_*(b_T); \mu_{b_*}, Q, \mu_Q) - S_{NP}^{h_1}(b_T, Q)]$$

$$\tilde{H}_1^{\perp(1)}(z, b_T; Q^2, \mu_Q) \sim \textcolor{blue}{H}_1^{\perp(1)}(z; \mu_{b_*}) \exp \left[-S_{pert}(b_*(b_T); \mu_{b_*}, Q, \mu_Q) - S_{NP}^{H_1^\perp}(b_T, Q) \right]$$




(Kang, et al. (2016))

$$p^\uparrow p \rightarrow hX$$




A_N is a *collinear* (twist-3) observable

$$d\Delta\sigma(S_T) \sim \underbrace{H_{QS} \otimes f_1 \otimes \textcolor{magenta}{F}_{FFT} \otimes D_1}_{\text{Qiu-Sterman term}} + \underbrace{H_F \otimes f_1 \otimes \mathbf{h}_1 \otimes (\mathbf{H}_1^{\perp(1)}, \tilde{\mathbf{H}})}_{\text{Fragmentation term}}$$



$$p^\uparrow p \rightarrow hX$$




A_N is a *collinear* (twist-3) observable

$$d\Delta\sigma(S_T) \sim \underbrace{H_{QS} \otimes f_1 \otimes \mathbf{F}_{FFT} \otimes D_1}_{\text{Qiu-Sterman term}}$$

$$+ \underbrace{H_F \otimes f_1 \otimes \mathbf{h}_1 \otimes (\mathbf{H}_1^{\perp(1)}, \tilde{\mathbf{H}})}_{\text{Fragmentation term}}$$


(Metz, DP (2012); Kanazawa, et al. (2014); Gumberg, et al. (2017); Cammarota, et al. (2020))







Simultaneous QCD Global Analysis of SSAs


Cammarota, Gamberg, Kang, Miller, DP, Prokudin, Rogers, Sato, PRD 102 (2020)



$$\chi^2/N_{\text{pts.}} = 520/517 = 1.01$$




JAM3D-20



Transversity

Sivers first
moment
(QS function)

Collins first
moment



Cammarota, Gamberg, Kang, Miller, DP, Prokudin, Rogers, Sato, PRD **102** (2020)

Only after a *simultaneous* QCD global analysis of SSAs does the phenomenological extraction of the tensor charges agree with lattice, *but still with large uncertainties*.





Impact Study on the Tensor Charge

Gamberg, Kang, DP, Prokudin, Sato, Seidl, PLB **816** (2021)

EIC Pseudo-data			
Observable	Reactions	CM Energy (\sqrt{S})	$N_{\text{pts.}}$
Collins (SIDIS)	$e + p^\uparrow \rightarrow e + \pi^\pm + X$	141 GeV	756 (π^+) 744 (π^-)
		63 GeV	634 (π^+) 619 (π^-)
		45 GeV	537 (π^+) 556 (π^-)
		29 GeV	464 (π^+) 453 (π^-)
		85 GeV	647 (π^+) 650 (π^-)
	$e + {}^3He^\uparrow \rightarrow e + \pi^\pm + X$	63 GeV	622 (π^+) 621 (π^-)
		29 GeV	461 (π^+) 459 (π^-)
		Total EIC $N_{\text{pts.}}$	8223


Assumed accumulated luminosities of 10 fb^{-1} , 70% polarization, conservatively accounted for detector smearing and acceptance effects

Gamberg, Kang, DP, Prokudin, Sato, Seidl, PLB 816 (2021)





EIC data on the Collins effect will significantly reduce the uncertainties in extractions of the transversity PDF (as well as the Collins FF)

Gamberg, Kang, DP, Prokudin, Sato, Seidl, PLB 816 (2021)




EIC data will allow phenomenological extractions of the tensor charge to become as precise as current lattice calculations. 3He data is especially important to decorrelate the extraction of δu and δd .

Gamberg, Kang, DP, Prokudin, Sato, Seidl, PLB 816 (2021)




SoLID at JLab covers a complimentary region at higher x and lower Q^2 with much greater luminosity – important to explore the effect of multiple measurements in different kinematic regions

Gamberg, Kang, DP, Prokudin, Sato, Seidl, PLB 816 (2021)



SoLID reduces the relative uncertainty in the down quark h_1 at higher x more than the EIC, and overall the relative uncertainties improve the most when data sets from both facilities are included

Gamberg, Kang, DP, Prokudin, Sato, Seidl, PLB 816 (2021)



SoLID at JLab will also provide important constraints on the tensor charges. The combined analysis with EIC data gives the most precise phenomenological determination of them.

Conclusions

- We have performed the first global analysis of SSAs in SIDIS, DY, e^+e^- annihilation, and proton-proton collisions and extracted a universal set of non-perturbative functions, showing a common origin of SSAs.
- First agreement with lattice QCD on the tensor charges of the nucleon was obtained, but still with large uncertainties.
- EIC data on the Collins effect will allow for phenomenological extractions of the tensor charge to be as precise as current lattice calculations. SoLID at JLab will also provide important constraints.
- In order to reduce bias and obtain the most accurate extraction of the tensor charge, one must have data from multiple future facilities (e.g., EIC and SoLID) that give the most kinematic coverage possible in x and Q^2 .



Back-up Slides

- We have performed the first global analysis of SSAs in SIDIS, Drell-Yan, e^+e^- annihilation, and proton-proton collisions and extracted a universal set of non-perturbative functions

$$h_1(x), F_{FT}(x, x), H_1^{\perp(1)}(z), \hat{H} \cancel{X} z) \quad \text{noise in the fit - need } A_{UT}^{\sin \phi_S}$$

along with the relevant transverse momentum widths for the Sivers, transversity, and Collins functions: $\langle k_T^2 \rangle_{f_{1T}^\perp}$, $\langle k_T^2 \rangle_{h_1}$, $\langle p_\perp^2 \rangle_{H_1^\perp}^{fav}$, $\langle p_\perp^2 \rangle_{H_1^\perp}^{unf}$

- We use a Gaussian ansatz: $F(x, k_T^2) \sim F(x) e^{-k_T^2 / \langle k_T^2 \rangle}$ where

$$F^q(x) = \frac{N_q x^{a_q} (1-x)^{b_q} (1 + \gamma_q x^{\alpha_q} (1-x)^{\beta_q})}{B[a_q+2, b_q+1] + \gamma_q B[a_q+\alpha_q+2, b_q+\beta_q+1]}$$

NB: $\{\gamma, \alpha, \beta\}$ only used for Collins function

- DGLAP-type evolution for the collinear functions analogous to Duke & Owens (1984): double-log Q^2 -dependent term explicitly added to the parameters

Observable	Reactions	Non-Perturbative Function(s)	$\chi^2/N_{\text{pts.}}$
$A_{\text{SIDIS}}^{\text{Siv}}$	$e + (p, d)^\uparrow \rightarrow e + (\pi^+, \pi^-, \pi^0) + X$	$f_{1T}^\perp(x, k_T^2)$	$150.0/126 = 1.19$
$A_{\text{SIDIS}}^{\text{Col}}$	$e + (p, d)^\uparrow \rightarrow e + (\pi^+, \pi^-, \pi^0) + X$	$h_1(x, k_T^2), H_1^\perp(z, z^2 p_\perp^2)$	$111.3/126 = 0.88$
$A_{\text{SIA}}^{\text{Col}}$	$e^+ + e^- \rightarrow \pi^+ \pi^- (\text{UC}, \text{UL}) + X$	$H_1^\perp(z, z^2 p_\perp^2)$	$154.5/176 = 0.88$
$A_{\text{DY}}^{\text{Siv}}$	$\pi^- + p^\uparrow \rightarrow \mu^+ \mu^- + X$	$f_{1T}^\perp(x, k_T^2)$	$5.96/12 = 0.50$
$A_{\text{DY}}^{\text{Siv}}$	$p^\uparrow + p \rightarrow (W^+, W^-, Z) + X$	$f_{1T}^\perp(x, k_T^2)$	$31.8/17 = 1.87$
A_N^h	$p^\uparrow + p \rightarrow (\pi^+, \pi^-, \pi^0) + X$	$h_1(x), F_{FT}(x, x) = \frac{1}{\pi} f_{1T}^{\perp(1)}(x), H_1^{\perp(1)}(z)$	$66.5/60 = 1.11$

- 18 observables and 6 non-perturbative functions (Sivers up/down; transversity up/down; Collins favored/unfavored)

**Test of
universality!**

- Broad kinematical coverage:

SIDIS: $x \lesssim 0.3$ $0.2 \lesssim z \lesssim 0.6$ $2 \lesssim Q^2 \lesssim 40 \text{ GeV}^2$



SIA: $0.2 \lesssim z \lesssim 0.8$ $Q^2 \approx 13 \text{ GeV}^2$ or 110 GeV^2

DY: $0.1 \lesssim x \lesssim 0.35$ $Q^2 \approx 30 \text{ GeV}^2$ or $(80 \text{ GeV})^2$

A_N^h : $0.2 \lesssim (x_{\min}, z_{\min}) \lesssim 0.7$ $1 \lesssim Q^2 \lesssim 13 \text{ GeV}^2$


Observable	Reactions	Non-Perturbative Function(s)	$\chi^2/N_{\text{pts.}}$
$A_{\text{SIDIS}}^{\text{Siv}}$	$e + (p, d)^\uparrow \rightarrow e + (\pi^+, \pi^-, \pi^0) + X$	$f_{1T}^\perp(x, k_T^2)$	$150.0/126 = 1.19$
$A_{\text{SIDIS}}^{\text{Col}}$	$e + (p, d)^\uparrow \rightarrow e + (\pi^+, \pi^-, \pi^0) + X$	$h_1(x, k_T^2), H_1^\perp(z, z^2 p_\perp^2)$	$111.3/126 = 0.88$
$A_{\text{SIA}}^{\text{Col}}$	$e^+ + e^- \rightarrow \pi^+ \pi^- (\text{UC}, \text{UL}) + X$	$H_1^\perp(z, z^2 p_\perp^2)$	$154.5/176 = 0.88$
$A_{\text{DY}}^{\text{Siv}}$	$\pi^- + p^\uparrow \rightarrow \mu^+ \mu^- + X$	$f_{1T}^\perp(x, k_T^2)$	$5.96/12 = 0.50$
$A_{\text{DY}}^{\text{Siv}}$	$p^\uparrow + p \rightarrow (W^+, W^-, Z) + X$	$f_{1T}^\perp(x, k_T^2)$	$31.8/17 = 1.87$
A_N^h	$p^\uparrow + p \rightarrow (\pi^+, \pi^-, \pi^0) + X$	$h_1(x), F_{FT}(x, x) = \frac{1}{\pi} f_{1T}^{\perp(1)}(x), H_1^{\perp(1)}(z)$	$66.5/60 = 1.11$

- Predictions of A_N using a fit of only TMD observables



Test of universality!

Gamberg, Kang, DP, Prokudin, Sato, Seidl, PLB **816** (2021)



Most impact on JAM20 will be at moderate to higher x , across multiple decades in Q^2 , and we again see the importance of the ^3He program