

22nd edition

# PANIC Lisbon Portugal

Particles and Nuclei International Conference



Measurements of Lepton-Jet Azimuthal Decorrelations  
and 1-Jettiness event shape  
at high  $Q^2$  in DIS  
with H1 experiment at HERA



Sookhyun Lee (University of Michigan, Ann Arbor)  
on behalf of the H1 collaboration

PANIC 2021

Sep 8, 2021

[sookhyun@umich.edu](mailto:sookhyun@umich.edu)





Sookhyun Lee (University of Michigan) PANIC 2021





# HERA (1992-2007)




- First (and so far the only)  $ep$  collider
- Electron and positron as well as polarized (~40%) runs.
- $E_e = 27.6 \text{ GeV}$ ,  $E_p = 920 \text{ GeV}$ ,  $\sqrt{s} = 319 \text{ GeV}$
- Total integrated luminosity of  $356.1 \text{ pb}^{-1}$

# Deep-inelastic Scattering (DIS) at HERA



- Legacy of HERA, ep collider studying DIS
  - Proton structure and parton shower
  - Diffractive physics,
  - Electroweak interactions
  - Beyond standard model (BSM) physics
  - Determination of strong coupling constant – single inclusive jets, dijets.




# Deep-inelastic Scattering (DIS) and jet physics at EIC




- Forthcoming EIC era
  - Proposed measurements involving jets in DIS such as jet substructure, global event shape, correlations are rich in physics such as
    - 3-dimensional description of nucleon and hadronization, flavor dependence, precision measurements for QCD and BSM.
  - Theoretical and experimental advancement
    - pQCD calculation at NNLO and beyond, pushing N3LL accuracy in resummation.
    - Theory frameworks dealing with multiscale problems such as transverse momentum dependent (TMD) and soft-collinear effective theory (SCET).
    - Grooming technique, ML, sensor technology of a few  $\mu\text{m}$ , etc.
  - Motivates us to revisit HERA data!

# H1 experiment at HERA

- Fully instrumented with
  - Trackers (silicon strip + drift chamber + multi-wire proportional chamber)
  - Calorimeter (Liquid-Argon + Lead- scintillating fiber, EM+ hadronic in both)
  - Solenoid, muon-chambers, etc.
- Particle flow algorithm used for charged/neutral particle reconstruction.
- Events triggered by requiring high-energetic cluster in LAr calorimeter
  - electron or hadron
  - > 99% efficient for  $y < 0.7$



# Neutral current DIS



- Neutral current DIS
  - $e p \rightarrow e' X$  ( $e$  :  $e+$  or  $e-$ )
- Kinematic variables

$$\begin{aligned}
 Q^2 &= -q^2 = -(k - k')^2 \\
 x &= \frac{Q^2}{2P \cdot q} \\
 y &= \frac{P \cdot q}{P \cdot k} \\
 s &= (P + k)^2 = \frac{Q^2}{x \cdot y}
 \end{aligned}$$



# DIS kinematics reconstruction

- Reconstruction methods

- $\Sigma$  method

$$y_\Sigma = \frac{\sum_{i \in HFS} (E - p_z)_i}{\sum_{i \in HFS} (E - p_z)_i + E_{e'}(1 - \cos\theta_{e'})}$$

$$Q_\Sigma^2 = \frac{E_{e'}^2 \sin\theta_{e'}^2}{1 - y} \quad \begin{matrix} HFS: \\ \text{hadronic} \\ \text{final state} \end{matrix}$$

- $I\Sigma$  method

$$y_{I\Sigma} = y_\Sigma, Q_{I\Sigma}^2 = Q_\Sigma^2$$

$$\chi_{I\Sigma} = \frac{E_{e'}}{E_p} \frac{\cos^2(\theta_{e'}/2)}{y_\Sigma}$$

→ Largely insensitive to initial state QED radiative effects


# Lepton-jet decorrelation

- Novel way of probing transverse motion of quarks in proton, i.e. quark TMD PDF.
- Exploit lepton-jet  $p_T$  balance in Born kinematics

Liu et al. PRL. 122, 192003 (2019)  
Gutierrez et al. PRL. 121, 162001 (2019)

Lepton-jet  $p_T$  imbalance

$$q_T = |\vec{k}_\perp^l + \vec{p}_\perp^{jet}|$$



# Lepton-jet decorrelation


- Noble way of probing transverse motion of quarks in proton, i.e. quark TMD PDF.
- Exploit lepton-jet  $p_T$  balance in Born kinematics

Liu et al. PRL. 122, 192003 (2019)  
 Gutierrez et al. PRL. 121, 162001 (2019)


$$\frac{d^5\sigma(\ell p \rightarrow \ell' J)}{dy_\ell d^2k_{\ell\perp} d^2q_\perp} = \sigma_0 \int d^2k_\perp d^2\lambda_\perp x f_q(x, k_\perp, \zeta_c, \mu_F) \\ \times H_{\text{TMD}}(Q, \mu_F) S_J(\lambda_\perp, \mu_F) \\ \times \delta^{(2)}(q_\perp - k_\perp - \lambda_\perp).$$

Lepton-jet  $p_T$  imbalance

$$q_T = |\vec{k}_\perp^l + \vec{p}_\perp^{\text{jet}}|$$



# Machine-learning based unfolding








- Unbinned multi-dimensional unfolding.
- Deep neural networks trained to learn detector effects.
- Weight assigned to each event in MC.
- Unfolded distributions are obtained by binning reweighted events.

PRL 124, 182001 (2020)

Omnifold



# Closure tests



# Jet transverse momentum and pseudorapidity



arXiv:2108.12376





- $p_T$  and  $\eta$  of jets reasonably well described by NNLO + NP.
- Rapgap of all MC best describes data.
- Overall shape difference seen for Cascade (TMD based) in both variables.

# Lepton-jet momentum imbalance and azimuthal angular distance



arXiv:2108.12376



- $q_T$  reasonably well described by NNLO in higher region and by TMD at lower region, similar pattern for  $\Delta\phi$ .
- Large overlap covered by data will help constrain matching between TMD and collinear pQCD frameworks.


# 1-jettiness global event shape

- 1-jettiness

- $\tau_1^b = \frac{2}{Q^2} \sum_{i \in X} \min\{q_J \cdot p_i, q_B \cdot p_i\},$

where  $q_J = xP + q$ ,  $q_B = xP$

- Global, Lorentz invariant and infrared-safe observable
- Sensitive to strong coupling constant  $\alpha_s$  and PDFs







Kang, Lee, Stewart, PRD 88 (2013) 054004

N2LL published, N3LL WIP

# In Breit Frame

- Breit frame
  - Only spatial component in exchanged boson momentum.
  - Direction of momentum reversed after boson-parton head-on collision, i.e. brick-wall frame.
- $\tau_Q \stackrel{\text{def}}{=} 1 - \frac{2}{Q} \sum_{i \in \mathcal{H}_C} p_{z,i}^{\text{Breit}}$   
JHEP 02 (2000) 001
- By momentum conservation
  - $\tau_Q = \tau_1^b$






# MC simulations

- Contributions of different particles to  $\tau_1^b$ 
  - Major contribution from tracks in central region and particles with  $E > 1$  GeV.
  - Dominated by well measured particles.
- Kinematic phase space:
  - $150 < Q^2 < 20,000$  GeV $^2$
  - $0.2 < y < 0.7$

# Single differential cross section

- Limits of  $\tau_1^b$ 
  - $\tau_1^b \rightarrow 0$  : two pencil-like jets
  - $\tau_1^b \gg 0$  : multi-jets
  - $\tau_1^b = 1$  : empty current hemisphere
- NNLO ( $\alpha_s^2$ ) prediction from NNLOJET + NP corrections
  - Description of far-tail region sensitive to fixed-order (FO) improved by higher order NNLO.
  - Near peak region, improvement by NP corrections (Pythia 8.3).




- Unfolded by bin-by-bin correction.
- QED radiative effects corrected.

# $Q^2$ and $y$ dependent cross sections



H1prelim-21-032


- $\tau_1^b$ -differential cross section in  $y - Q^2$  space.
- Jets increasingly collimated with  $Q^2$ .
- NP effects (shift of peak position) more prominent in low  $Q^2$ .
- As  $y$  increases  $\tau_1^b = 1$  peak enhanced.



# $Q^2$ and $y$ dependent cross sections

H1prelim-21-032

- NNLO pQCD predictions
  - High  $y$  - high  $Q^2$  regions well described by NNLO.
  - NP corrections sizable at low  $Q^2$ .
  - Small scale uncertainty.
- Predictions with N3LL accuracy in SCET framework available soon.
  - Extended range in  $\tau_1^b$ , e.g. peak to far region with a few percent level theory uncertainty (a percent level constraining power on  $\alpha_s$ ) expected in high  $Q$  - high  $y$  region.







# Summary and Outlook

- First measurements of **lepton jet momentum imbalance and azimuthal decorrelation** in NC DIS presented as a new way to constrain TMD PDFs and their evolution.
- TMD calculation describes low  $q_T$  region of measured data well and pQCD collinear prediction does large  $q_T$  region. Large overlap region covered by data can constrain matching between TMD and collinear frameworks.
- First measurement of **1-jettiness event shape observable** in NC DIS was presented as a new way to greatly improve precision of PDF and  $\alpha_s$  determination.
- NNLO ( $\alpha_s^2$ ) fixed order predictions provide good description in the region of validity, but hadronization corrections are sizable.
- N3LL and NNLO+PS predictions will be compared with data.
- Sensitivity to  $\alpha_s$  and PDFs needs to be explored.
- Data will become useful for improving description of less inclusive DIS MC generators.

# backup

# Single differential cross section



- Djangoh 1.4 : color-dipole model
- Rapgap 3.1 : ME + parton shower
- Dire: Dipole-like shower + inclusive NLO DGLAP corrections.

# Q<sup>2</sup> and x dependent cross sections

• ...



# Q<sup>2</sup> and x dependent cross sections

- ...