

# Proton Decay Amplitudes with Physical Chirally-Symmetric Quarks

Taku Izubuchi, Amarjit Soni,  
**Sergey Syritsyn**, Jun-Sik Yoo



**PANIC 2021**

*Lisbon, September 8, 2021*

- Proton decay basics

  - Experimental lifetime limits & outlook*

  - Motivation and theory status*

  - Effective nucleon decay operators and matrix elements*

- Need for lattice calculations

  - Past calculations and model uncertainty*

  - Summary of the present calculation*

- Lattice calculation and analysis

  - Hadron masses and energies*

  - Extraction of matrix elements*

  - Operator renormalization*

  - Momentum & continuum extrapolations*


- Results

  - Comparison to earlier calculations*

  - Nucleon annihilation amplitudes*

  - Conclusions*

# Proton Stability: Status and Outlook




- Current limits  $\tau(p \rightarrow e^+ \pi^0) \gtrsim 1.6 \cdot 10^{34}$  ,  $\tau(p \rightarrow \bar{\nu} K^+) \gtrsim 5.9 \cdot 10^{33}$  [Super-K]
- Expect x10 improvement on lifetime limit from Hyper-K and DUNE
- Better sensitivity to  $p \rightarrow \bar{\nu} K^+$  that affects supersymmetric GUT models

# Motivation and Theory Status


- Proton lifetime tests *baryon number conservation*
  - *accidental "symmetry" of SM*, violated by sphalerons
  - has to be violated for baryogenesis [A.Sakharov'67]
- Missing piece of Grand-Unified Theories
- Probes scales inaccessible to colliders: Limits on GUT, extra dim., etc
- Limits on stability of nuclear matter

[Sakai, Yanagida '82; Weinberg '82]



## ordinary GUT

- min. $SU(5)$  ruled out by  $\tau(p \rightarrow e^+ \pi^0)$
- $SO(10)$  probed by next-gen exp.



## supersymmetric GUT

- min.SUSY- $SU(5)$  ruled out by  $\tau(p \rightarrow \bar{\nu} K^+)$
- SUSY- $SO(10)$  probed by next-gen exp.


# How Nucleon Structure Affects GUT Limits

Effective interaction

$$\mathcal{L}_{\text{eff}} = \sum_I C_I \mathcal{O}_I + \text{h.c.}$$

$$\mathcal{O}_I = \epsilon^{abc} (\bar{q}_1^{aC} P_{\chi_I} q_2^b) (\bar{\ell}^C P_{\chi'_I} q_3^c) = \bar{\ell}_\alpha^C \mathcal{O}_{I,\alpha}^{3q}$$

$$q_{1,2,3} \in \{u, d, s\}, \quad P_{\chi_I^{(\prime)}} = \frac{1 \pm \gamma_5}{2}$$



Decay width  $p \rightarrow \Pi \bar{\ell}$  ( $\Pi = \pi, K, \eta$ ;  $m(\bar{\ell}) \ll m_N$ )

$$\Gamma(p \rightarrow \Pi \bar{\ell}) = \frac{m_N}{32\pi} \left[ 1 - \left( \frac{m_\Pi}{m_N} \right)^2 \right]^2 \left| \sum_I C_I W_{\bar{\ell}}^I \right|^2$$

Decay matrix elements  $(W_{0,1})_I$

[S.Aoki et al, PRD62:014506 (200)]

$$\langle \bar{\ell}(q) \Pi(p) | \mathcal{O}^{x'} | N(k) \rangle = \bar{v}_{\ell\alpha}^C(q) P_{\chi'} \left[ W_0(-q^2) - \frac{i \not{q}}{m_N} W_1(-q^2) \right] u_N(k)$$

$$\text{and } W_{\bar{\ell}} = \left[ W_0 + W_1 \cdot O(m_{\bar{\ell}}/m_N) \right]_{q^2=m_{\bar{\ell}}^2}$$

*nonperturbative QCD ;  
require lattice calculation*

negligible for  $e^+$  ;  
 $\approx 10\%$  for  $\mu^+$

# Nucleon Decay Matrix Elements

Nonperturbative matrix elements [form factors]

$$\langle \Pi(k - q) | \mathcal{O}_\alpha^{3q} | N(k) \rangle = \left[ P_{\chi'} \left( W_0^\mathcal{O} - \frac{i \not{q}}{m_N} W_1^\mathcal{O} \right) u_N(k) \right]_\alpha$$

[S.Aoki et al, PRD62:014506 (200)]

Two methods to calculate  $W_{0,1}$

- Direct calculation on lattice
- Low-energy theory (soft-pion thm.)  
requires annihilation amplitude  $\langle \text{vac} | \mathcal{O}^{3q} | N \rangle$   
(also needed for  $p \rightarrow 3\bar{\ell}$  decays)


*Order-of-magnitude estimate:*

$$\langle \text{vac} | \mathcal{O}^{3q} | N \rangle \sim \rho_q^{3/2} \sqrt{V_N} \sim \frac{1}{V_N} \approx 0.004 \text{ GeV}^3$$

$$\langle \Pi | \mathcal{O}^{3q} | N \rangle \sim \langle \text{vac} | \mathcal{O}^{3q} | N \rangle / f_\pi \approx 0.03 \text{ GeV}^2$$


[Martin, Stavenga '12]

Suppression of  $\langle \text{vac} | \mathcal{O}^{3q} | N \rangle$   
in Chiral Bag model  
due skyrmion topology



*Alternative explanation for  
observed proton stability?*

lattice calculations  
with  $m_\pi \approx 330 \text{ MeV}$



# This Work: Lattice Setup

- Two ensembles: [32ID]  $32^3 \times 64 (a=0.14 \text{ fm})$  and [24ID]  $24^3 \times 64 (a=0.20 \text{ fm})$
- Iwasaki gauge action+ Dislocation-supp. det.ratio (DSDR)
- $N_f = 2+1$  Chirally-symmetric (Mobius-)Domain Wall fermion action with physical light and strange quark masses
- Multigrid deflation of z-Mobius operator + AMA
- "Direct" ( $p \rightarrow \pi, K$  matrix elements) and "Indirect" ( $p \rightarrow \text{vacuum} + \text{ChPT}$ )
- Nonperturbative renormalization
- Two state-fit analysis of  $\pi, K, N$  spectrum and  $p \rightarrow \pi, K$  matrix elements
- $a^2$  Continuum extrapolation





	24ID $24^3 \times 64$	32ID $32^3 \times 64$
$\beta$	1.633	1.75
$a, \text{ fm}$	0.20	0.14
$a^{-1}, \text{ GeV}$	1.02	1.37
$m_\pi L$	3.4	3.3
$N_{conf}$	134	94
$N_{samp}$	4288	3008

- three kinematic ( $Q^2$ ) points to interpolate matrix elements to decay kinematic  $Q^2 = -(m_{\bar{\ell}})^2$


$\Pi$	$\vec{n}_\Pi$	$\vec{n}_N$	$Q^2 (\text{GeV}^2)$	
			(24c)	(32c)
$\pi$	[1 1 1]	[0 0 0]	0.010	-0.012
	[1 1 1]	[0 1 0]	0.113	0.095
	[0 0 2]	[0 0 0]	-0.116	-0.140
$K$	[0 1 1]	[0 0 0]	-0.034	-0.042
	[0 1 1]	[0 1 0]	0.058	0.056
	[0 0 1]	[0 0 0]	0.075	0.074




# Proton and Meson Spectrum



**pion**



**kaon**



**nucleon**


- 24ID ensemble ( $a=0.20$  fm)
- Two-state fits + priors from large- $t_{\min}$  one-state fits



# Extraction of Matrix Elements

32ID

$W_0$



- Two-state fits with energies fixed from spectrum fits: *no signs of excited-state sys.errors*

# Nonperturbative Renormalization




- symmetry-allowed mixing

	$\mathcal{S} = -1$	$\mathcal{S} = +1$
$\mathcal{P} = -1$	$SS, PP, AA$	$VV, TT$
$\mathcal{P} = +1$	$SP, PS, AV$	$VA, TQ$

- symmMOM* scheme :  $p+q+r=0$ ,  $p^2=q^2=r^2=\mu^2$


$$Z_{IK}^{3q}(\mu) \text{Proj}_J [\langle \bar{q}_1(p) \bar{q}_2(q) \bar{q}_3(r) \mathcal{O}_K^{3q} \rangle_{\text{amp}}] = \delta_{IJ}$$

- symmMOM*( $p$ ) $\rightarrow$ MSbar(2 GeV)  
perturbative conversion at  $O(\alpha^3)$   
[J.Gracey, JHEP09:052 (2012)]





- chiral symmetry suppresses mixing of  $L \rightleftharpoons R$  fields & operators*

# Momentum and Continuum Extrapolation



- linear momentum extrapolation  $Q^2 \rightarrow m_e^2, m_\mu^2$  to the decay kinematics
- Continuum extrapolation  $A(a^2) \sim (A_0 + A_2 a^2)$  ; *conservative* sys.error =  $|A_0 - A_{[a=0.14\text{fm}]}|$

# Proton Annihilation Amplitudes



$$\langle \text{vac} | \epsilon^{abc} (\bar{u}^{aC} d^b)_R u_L^c | N \rangle = \alpha P_L U_N$$


$$\langle \text{vac} | \epsilon^{abc} (\bar{u}^{aC} d^b)_L u_L^c | N \rangle = \beta P_L U_N$$

- connected to  $\langle \pi/K | O^{3q} | N \rangle$  by soft-pion theorem


- $(\alpha + \beta) = 0$  [within errorbars] implying

$$\epsilon^{abc} (\bar{u}^{aC} d^b) \gamma_5 u^c | N \rangle \stackrel{?}{\approx} 0$$

parity                      (−)      (−)      (+)



# Comparison to Previous Work



- **New results:**  
(stat+sys) precision  $\sim 10\text{-}20\%$
- No FVE study,  $\text{mpi} \cdot L \sim 3.4$
- physical-point results agree with  
prev. calculations at  $m_\pi \gtrsim 300$  MeV  
[S.Aoki et al (2000)]  
[Y.Aoki et al (2006)]  
[Y.Aoki et al (2013)]

*NO suppression of nucleon decay  
due to chiral skyrmion topology*

# Summary & Conclusions

- *Proton decay amplitudes at the physical point with chiral symmetric quarks and continuum extrapolation*
- *No topological suppression of nucleon decay found; limits on Grand-Unified Theories **stand***
- *Sys+Stat. precision  $O(10-20\%)$  ; may be improved with more statistics, finer lattice spacing, finite-volume study*

# BACKUP