

# Phenomenological study of quarkonium suppression and the impact of the energy gap between singlets and octets

Miguel A. Escobedo

Instituto Galego de Física de Altas Enerxías  
Universidade de Santiago de Compostela

September 8, 2021



FONDO EUROPEO DE DESENVOLVEMENTO REXIONAL  
"Unha maneira de facer Europa"



EXCELENCIA  
MARÍA  
DE MAEZTU



Xacobeo 2021

# References

- Blaizot and Escobedo, Quantum and classical dynamics of heavy quarks in a quark-gluon plasma, JHEP 06 (2018) 034.
- Blaizot and Escobedo, Approach to equilibrium of a quarkonium in a quark-gluon plasma, Phys.Rev.D 98 (2018) 7, 074007.
- Blaizot and Escobedo, Phenomenological study of quarkonium suppression and the impact of the energy gap between singlets and octets, arXiv:2106.15371

1 Introduction

2 Derivation

3 Phenomenological application

4 Conclusions

# Heavy quarkonium in heavy-ion collisions

- Heavy quarkonium is a bound state of heavy quarks, whose mass is larger than  $\Lambda_{QCD}$ .

# Heavy quarkonium in heavy-ion collisions

- Heavy quarkonium is a bound state of heavy quarks, whose mass is larger than  $\Lambda_{QCD}$ .
- Heavy quarks can only be created at the beginning of the collision. It is a hard process.

# Heavy quarkonium in heavy-ion collisions

- Heavy quarkonium is a bound state of heavy quarks, whose mass is larger than  $\Lambda_{QCD}$ .
- Heavy quarks can only be created at the beginning of the collision. It is a hard process.
- However, the existence of a medium changes the probability that a bound state is formed and its lifetime.

# Heavy quarkonium in heavy-ion collisions

- Heavy quarkonium is a bound state of heavy quarks, whose mass is larger than  $\Lambda_{QCD}$ .
- Heavy quarks can only be created at the beginning of the collision. It is a hard process.
- However, the existence of a medium changes the probability that a bound state is formed and its lifetime.
- Measuring  $R_{AA}$ , the ratio of quarkonium states measured in heavy-ion collisions divided by the naive extrapolation of pp data, we can extract information about the medium.

# Quarkonium as an Open quantum system

- We consider a *universe* consisting in heavy quarks (system) in a medium of light quarks and gluons (environment). The density matrix  $\rho(S, E)$  describes the state of the universe. Its evolution is unitary.




# Quarkonium as an Open quantum system

- We consider a *universe* consisting in heavy quarks (system) in a medium of light quarks and gluons (environment). The density matrix  $\rho(S, E)$  describes the state of the universe. Its evolution is unitary.
- We define a reduced density matrix integrating out the environment degrees of freedom  $\rho_S(S) = \text{Tr}_E(\rho(S, E))$ . We study the evolution of  $\rho_S$  that, in general, is not unitary.

# Quarkonium as an Open quantum system

- We consider a *universe* consisting in heavy quarks (system) in a medium of light quarks and gluons (environment). The density matrix  $\rho(S, E)$  describes the state of the universe. Its evolution is unitary.
- We define a reduced density matrix integrating out the environment degrees of freedom ( $\rho_S(S) = \text{Tr}_E(\rho(S, E))$ ). We study the evolution of  $\rho_S$  that, in general, is not unitary.



# The master equation

We call master equation the equation that describes the evolution of the reduced density matrix.

$$\frac{d\rho}{dt} = -i[H, \rho(t)] + \mathcal{F}(t, \rho(t))$$

# The master equation

We call master equation the equation that describes the evolution of the reduced density matrix.

$$\frac{d\rho}{dt} = -i[H, \rho(t)] + \mathcal{F}(t, \rho(t))$$

- We can recover the Schrödinger equation and the Boltzmann equation as limits of the master equation in specific regimes.

# The master equation

We call master equation the equation that describes the evolution of the reduced density matrix.

$$\frac{d\rho}{dt} = -i[H, \rho(t)] + \mathcal{F}(t, \rho(t))$$

- We can recover the Schrödinger equation and the Boltzmann equation as limits of the master equation in specific regimes.
- We need to derive the master equation from QCD. This has been done in:
  - ▶ Perturbation theory. Akamatsu (2015,2020), Blaizot and Escobedo (2017,2018).
  - ▶ Potential non-relativistic QCD (pNRQCD) in the  $\frac{1}{r} \gg T$  regime. Brambilla et al. (2016,2017).

# History and motivation

- The QED approximation was studied by Blaizot et al. They found that the late time behaviour was well described by a Langevin equation.

# History and motivation

- The QED approximation was studied by Blaizot et al. They found that the late time behaviour was well described by a Langevin equation.
- Prof. Blaizot and me studied the QCD generalization. We found Langevin-like equations, but their applicability region is very narrow.

# History and motivation

- The QED approximation was studied by Blaizot et al. They found that the late time behaviour was well described by a Langevin equation.
- Prof. Blaizot and me studied the QCD generalization. We found Langevin-like equations, but their applicability region is very narrow.
- We understood why. In QED, the Langevin equation minimizes the classical free energy if it exists. This happens even if in the derivation approximations only valid when  $T \gg E$  are used.



# History and motivation

- The QED approximation was studied by Blaizot et al. They found that the late time behaviour was well described by a Langevin equation.
- Prof. Blaizot and me studied the QCD generalization. We found Langevin-like equations, but their applicability region is very narrow.
- We understood why. In QED, the Langevin equation minimizes the classical free energy if it exists. This happens even if in the derivation approximations only valid when  $T \gg E$  are used.
- These nice properties are lost in QCD. Crucial role of the energy gap between singlets and octets.

# History and motivation

- The QED approximation was studied by Blaizot et al. They found that the late time behaviour was well described by a Langevin equation.
- Prof. Blaizot and me studied the QCD generalization. We found Langevin-like equations, but their applicability region is very narrow.
- We understood why. In QED, the Langevin equation minimizes the classical free energy if it exists. This happens even if in the derivation approximations only valid when  $T \gg E$  are used.
- These nice properties are lost in QCD. Crucial role of the energy gap between singlets and octets.
- We wish to understand what is the phenomenological impact of taking into account this energy gap.

# Plan

1 Introduction


**2 Derivation**

3 Phenomenological application

4 Conclusions

# The evolution of the density matrix

Diagrams that connect whatever state at time  $t$  with a singlet at time  $t + dt$ .



$U_{s,o}$  is the LO time evolution operator. The quantum master equation can be written schematically as

$$\begin{aligned} \frac{d\mathcal{D}_Q}{dt} + i[H_Q, \mathcal{D}_Q(t)] = & \\ - \int_{\mathbf{x}\mathbf{x}'} \int_0^{t-t_0} d\tau [n_{\mathbf{x}}^A, U_Q(\tau)n_{\mathbf{x}'}^A \mathcal{D}_Q(t-\tau)U_Q^\dagger(\tau)] \Delta^>(\tau; \mathbf{x} - \mathbf{x}') & \\ - \int_{\mathbf{x}\mathbf{x}'} \int_0^{t-t_0} d\tau [U_Q(\tau)\mathcal{D}_Q(t-\tau)n_{\mathbf{x}'}^A U_Q^\dagger(\tau), n_{\mathbf{x}}^A] \Delta^<(\tau; \mathbf{x} - \mathbf{x}'), & \end{aligned}$$

where we have set  $t - t' = \tau$ .

## $U_{s,o}$ and energy conservation

The role of  $U_{s,o}$  is to inform the master equation of the size and the sign of the energy different between the state at  $t'$  and the state at  $t$ .

- If we make the approximation  $U_{s,o} = 1$  all transitions are equally likely.
- If  $U_{s,o}$  is not approximated there are some transitions which are more likely because they liberate energy while others are less likely because they need to absorb energy.
- $\frac{\Gamma(A \rightarrow B)}{\Gamma(B \rightarrow A)}$  is fixed by fluctuation-dissipation theorem.
- This is very important to understand QCD. An octet to singlet transition is always energetically favourable while the opposite happens for singlet to octet.

# The classical limit

The relation between the density matrix in the Schrödinger picture and in interaction picture is the following

$$D(t) = e^{-iHt} D_I(t) e^{iHt}$$

In the large time limit non-diagonal elements are suppressed by the fast oscillations

$$\mathcal{D}(t) = \sum_{nm} e^{-i(E_n - E_m)t} |n\rangle \langle n| \mathcal{D}_I(t) |m\rangle \langle m| \sim p_n(t) |n\rangle \langle n|$$

Evolution equation for the density matrix  $\rightarrow$  rate equation

## Rate equations

In a finite volume (octet levels are discrete)

$$\frac{dp_n^s}{dt} = 4g^2 C_F \sum_m \left( p_m^o - p_n^s e^{-\frac{E_m^o - E_n^s}{T}} \right) \int_{\mathbf{q}} \Gamma^>(E_m^o - E_n^s, \mathbf{q}) |\langle n^s | \mathcal{S}_{\mathbf{q},\mathbf{r}} | m^o \rangle|^2$$

and

$$\begin{aligned} \frac{dp_m^o}{dt} = & -\frac{2g^2}{N_c} \sum_n \left( p_m^o - p_n^s e^{-\frac{E_m^o - E_n^s}{T}} \right) \int_{\mathbf{q}} \Gamma^>(E_m^o - E_n^s, \mathbf{q}) |\langle n^s | \mathcal{S}_{\mathbf{q},\mathbf{r}} | m^o \rangle|^2 \\ & -\frac{g^2(N_c^2 - 4)}{N_c} \sum_k \left( p_m^o - p_k^o e^{-\frac{E_m^o - E_k^o}{T}} \right) \int_{\mathbf{q}} \Gamma^>(E_m^o - E_k^o, \mathbf{q}) |\langle m^o | \mathcal{S}_{\mathbf{q},\mathbf{r}} | k^o \rangle|^2 \\ & -g^2 N_c \sum_k \left( p_m^o - p_k^o e^{-\frac{E_m^o - E_k^o}{T}} \right) \int_{\mathbf{q}} \Gamma^>(E_m^o - E_k^o, \mathbf{q}) |\langle m^o | \mathcal{C}_{\mathbf{q},\mathbf{r}} | k^o \rangle|^2. \end{aligned}$$

# Free energy minimization

$$\begin{aligned}
 \frac{dF}{dt} = & -4g^2 C_F T \sum_{nm} \log \left( \frac{p_m^o}{p_n^s e^{-\frac{E_m^o - E_n^s}{T}}} \right) \left( p_m^o - p_n^s e^{-\frac{E_m^o - E_n^s}{T}} \right) \\
 & \times \int_{\mathbf{q}} \Delta^>(E_m^o - E_n^s, \mathbf{q}) |\langle n^s | \mathcal{S}_r | m^o \rangle|^2 \\
 & - \frac{g^2 T (N_c^2 - 4)}{2N_c} \sum_{mk} \log \left( \frac{p_m^o}{p_k^o e^{-\frac{E_m^o - E_k^o}{T}}} \right) \left( p_m^o - p_k^o e^{-\frac{E_m^o - E_k^o}{T}} \right) \\
 & \times \int_{\mathbf{q}} \Delta^>(E_m^o - E_k^o, \mathbf{q}) |\langle m^o | \mathcal{S}_r | k^o \rangle|^2 \\
 & - \frac{g^2 T N_c}{2} \sum_{mk} \log \left( \frac{p_m^o}{p_k^o e^{-\frac{E_m^o - E_k^o}{T}}} \right) \left( p_m^o - p_k^o e^{-\frac{E_m^o - E_k^o}{T}} \right) \\
 & \times \int_{\mathbf{q}} \Delta^>(E_m^o - E_k^o, \mathbf{q}) |\langle m^o | \mathcal{C}_{\mathbf{q},r} | k^o \rangle|^2 \leq 0.
 \end{aligned}$$



# Summary

- We can justify a rate equation if  $\Delta E$  (binding energy difference) is much bigger than the decay width.
- This can be justified (in some cases) for the singlet to octet transition. Not for octet to octet at a large volume (continuous spectrum).
- However, octet to octet transitions is similar to the QED case, which is reasonably described with a Langevin equation.

**Conclusion:** We can combine a rate equation for singlet to octet transitions with a Langevin equation for octet to octet.

## Our model

- In the large  $N_c$  limit the octet potential is zero.
- As an illustration we consider only one singlet eigenvalue (the 1S).  
This is not too bad since the 2S melts at a much smaller temperature.

$$\frac{dp^s}{dt} = g^2 C_F \int_{\mathbf{p}} \left( p_{\mathbf{p}}^o - p^s e^{-\frac{E_{\mathbf{p}}^o - E^s}{T}} \right) \int_{\mathbf{q}} \Delta^>(\omega_{\mathbf{p}}^o - E^s, \mathbf{q}) |\langle s | \mathcal{S}_{\mathbf{q}, \hat{r}} | o, \mathbf{p} \rangle|^2,$$

and

$$\begin{aligned} \frac{\partial p_{\mathbf{p}}^o}{\partial t} - \gamma \nabla(\mathbf{p} p_{\mathbf{p}}^o) - \frac{T \gamma M}{2} \Delta^2 p_{\mathbf{p}}^o = \\ - \frac{g^2}{2N_c} \frac{1}{\Omega} \left( p_{\mathbf{p}}^o - p^s e^{-\frac{E_{\mathbf{p}}^o - E^s}{T}} \right) \int_{\mathbf{q}} \Delta^>(\omega_{\mathbf{p}}^o - E^s, \mathbf{q}) |\langle s | \mathcal{S}_{\mathbf{q}, \hat{r}} | o, \mathbf{p} \rangle|^2, \end{aligned}$$

## Our model. Results

	$\Omega = 1 \text{ fm}^3$			$\Omega = 100 \text{ fm}^3$		
	5 fm/c	100 fm/c	eq.	5 fm/c	100 fm/c	eq.
$T = 200 \text{ MeV}$	0.86	0.136	0.0814	0.85	0.0438	0.00089
$T = 400 \text{ MeV}$	0.39	0.0515	0.0175	0.36	0.0002	0.00018

Table:  $p^s$  as obtained by solving eqs. in previous slide

### Remarks

- The gap  $E_p^o - E^s$  suppresses the singlet decay width, helping to make the results compatible with experimental observations.
- $E^s$  depends on the real part of the potential. Screening reduces the binding energy and this increases the decay width.
- The volume of the medium  $\Omega$  suppresses the decay of octets into bound singlets. Also suppressed in the large- $N_c$  limit.

# Plan

- 1 Introduction
- 2 Derivation
- 3 Phenomenological application**
- 4 Conclusions

# Outline

In this section we are going to study the impact of considering the finite gap between singlets and octets in phenomenological applications. We are going to consider two different scenarios:

- HTL inspired case.
- Lattice inspired case, based on the static potential data of Lafferty and Rothkopf (2020).

We will compute  $R_{AA}$  for these two cases assuming Bjorken evolution of the medium.

# Procedure

We solve the equation

$$\frac{dp_n}{dt} = -\Gamma_n p_n$$

and we compare the results that we get including (or not) the finite gap in the computation of  $\Gamma_n$ . For several temperatures:

- 1 We solve the Schrödinger equation to obtain the binding energy and the wave function.
- 2 We use the binding energy and the wave function to compute the decay width.

Once we have done this for several temperatures, we fit the decay width as a function of the temperature.

## The decay width

$$\Gamma_s = 8\pi\alpha_s C_F \int_{\mathbf{p}} e^{-\frac{E_{\mathbf{p}}^o - E^s}{T}} \int_{\mathbf{q}} \Delta^>(E_{\mathbf{p}}^o - E^s, \mathbf{q}) |\langle s | \sin\left(\frac{\mathbf{q} \cdot \hat{\mathbf{r}}}{2}\right) | o, \mathbf{p} \rangle|^2$$

where  $\Delta^>$  is the HTL  $A_0$  propagator in the Coulomb gauge. If we neglect the energy gap between singlets and octets we obtain

$$\Gamma_s^0 = 8\pi\alpha_s C_F \int_{\mathbf{q}} \Delta^>(0, \mathbf{q}) \langle s | \sin^2\left(\frac{\mathbf{q} \cdot \hat{\mathbf{r}}}{2}\right) | s \rangle$$

Note that this is equal to the expectation value of the imaginary part of Laine potential.

# Perturbative scenario

## Potential

$$V_s(r) = -\frac{C_F\alpha_s(\mu_r)e^{-m_D(T,\mu_T)r}}{r}$$


where  $\mu_r$  and  $\mu_T$  are (independent) subtraction points.  $\mu_r$  is either  $\frac{1}{2a_0}$ ,  $\frac{1}{a_0}$  or  $\frac{2}{a_0}$ , with  $a_0$  the Bohr radius.  $\mu_T$  is either  $\pi T$ ,  $2\pi T$  and  $4\pi T$ .

$$\frac{1}{a_0} = \frac{MC_F\alpha_s(1/a_0)}{2}, \quad a_0 = 0.149 \text{ fm}$$



# Perturbative scenario


## The Debye mass



$$m_D(T) = \sqrt{\frac{4\pi\alpha_s(\mu_T)(N_c + T_F N_F)}{3}} T$$

# Perturbative scenario

Dimensionless quantities for  $\Upsilon(1S)$



For  $\Upsilon(2S)$  the condition  $E \gg \Gamma$  is not fulfilled for this scenario.

# Perturbative scenario


$\Gamma$ , fit function

$$\Gamma \sim aT + bT^2$$

This function fits well the values of  $\Gamma$  for different temperatures. We computed the values of  $\Gamma$  from  $T = 200 \text{ MeV}$  to  $550 \text{ MeV}$  in intervals of  $50 \text{ MeV}$


# Perturbative scenario

$\Gamma$  of  $\Upsilon(1S)$




# Perturbative scenario

$\Gamma$  of  $\Upsilon(1S)$




# Perturbative scenario

$\Gamma$  of  $\Upsilon(1S)$ , without gap



# Perturbative scenario

$\Gamma$  of  $\Upsilon(1S)$ , without gap



# Perturbative scenario

## Survival probability

$$S_{(1S)}(\mathbf{b}, \mathbf{s}) = e^{-1.5aT_0(\mathbf{b}, \mathbf{s})t_0 \left( \left( \frac{T_0(\mathbf{b}, \mathbf{s})}{T_f} \right)^2 - 1 \right) - 3bT_0(\mathbf{b}, \mathbf{s})^2 t_0 \left( \frac{T_0(\mathbf{b}, \mathbf{s})}{T_f} - 1 \right)}$$

Survival probability of a  $\Upsilon(1S)$  state with  $\Gamma = aT + bT^2$  in a medium that follows Bjorken evolution up to a temperature  $T_f = 200 \text{ MeV}$ .



# Lattice scenario

## The potential

$$V_s(r) = -\frac{\alpha e^{-m_D r}}{r} - \sigma r e^{-m_D r} \left(1 + \frac{2}{m_D r}\right)$$

- Potential used in Lafferty and Rothkopf (2020) to fit lattice data.  $m_D$  is a fit parameter.

# Lattice scenario

## The potential

$$V_s(r) = -\frac{\alpha e^{-m_D r}}{r} - \sigma r e^{-m_D r} \left( 1 + \frac{2}{m_D r} \right)$$

- Potential used in Lafferty and Rothkopf (2020) to fit lattice data.  $m_D$  is a fit parameter.
- We observe that the biggest source of uncertainty from this fit comes from the parameter  $m_D$ .

# Lattice scenario


## The potential

$$V_s(r) = -\frac{\alpha e^{-m_D r}}{r} - \sigma r e^{-m_D r} \left(1 + \frac{2}{m_D r}\right)$$

- Potential used in Lafferty and Rothkopf (2020) to fit lattice data.  $m_D$  is a fit parameter.
- We observe that the biggest source of uncertainty from this fit comes from the parameter  $m_D$ .
- We use this potential for temperatures such that the energy gap is smaller  $2M_B - M_{\Upsilon(1S)}$ . If this is not the case we assume that there are no thermal modifications.


# Lattice scenario

$\Upsilon(1S)$  binding energy



# Lattice scenario

$\Upsilon(2S)$  binding energy



# Lattice scenario

Fit function

For  $\Upsilon(1S)$


$$\Gamma_s \sim \tilde{a} T e^{-\frac{\tilde{b}}{T}}$$

For  $\Upsilon(2S)$

$$\Gamma \sim aT + bT^2$$


# Lattice scenario

$\Upsilon(1S)$  decay width



# Lattice scenario


$\Upsilon(2S)$  decay width






# Lattice scenario

$\Upsilon(1S)$  decay width, without gap



# Lattice scenario

$\Upsilon(2S)$  decay width, without gap



# Lattice scenario

## Survival probability

For  $\Upsilon(1S)$

$$S_{(1S)}(\mathbf{b}, \mathbf{s}) = e^{-\frac{3\tilde{a}T_0(\mathbf{b}, \mathbf{s})^3 t_0}{\tilde{b}^2} \left( e^{-\frac{\tilde{b}}{T_0(\mathbf{b}, \mathbf{s})}} \left( 1 + \frac{\tilde{b}}{T_0(\mathbf{b}, \mathbf{s})} \right) - e^{-\frac{\tilde{b}}{T_f}} \left( 1 + \frac{\tilde{b}}{T_f} \right) \right)}$$

For  $\Upsilon(2S)$

$$S_{(2S)}(\mathbf{b}, \mathbf{s}) = e^{-1.5aT_0(\mathbf{b}, \mathbf{s})t_0 \left( \left( \frac{T_0(\mathbf{b}, \mathbf{s})}{T_f} \right)^2 - 1 \right) - 3bT_0(\mathbf{b}, \mathbf{s})^2 t_0 \left( \frac{T_0(\mathbf{b}, \mathbf{s})}{T_f} - 1 \right)}$$

In both case  $T_f = 180 \text{ fm}$

# The temperature of the medium and its evolution

$$T_0(\mathbf{b}, \mathbf{s}) = T_0(0, 0) \left( \frac{T_A(\mathbf{s}, 0) \left[ 1 - \left( 1 - \frac{\sigma T_A(\mathbf{s}-\mathbf{b}, 0)}{A} \right)^A \right]}{T_A(0, 0) \left[ 1 - \left( 1 - \frac{\sigma T_A(0, 0)}{A} \right)^A \right]} \right)^{1/4}$$

We take  $T_0(0, 0) = 475 \times 1.05$  for  $\sqrt{s} = 5.02$  TeV collisions at LHC.

We assume Bjorken flow with  $t_0 = 0.6$  fm


$$T(t, \mathbf{b}, \mathbf{s}) = T_0(\mathbf{b}, \mathbf{s}) \left( \frac{t_0}{t} \right)^{\frac{1}{3}}$$

## Computation of $R_{AA}$


$$R_{AA}(\mathbf{b}) = \frac{\int d^2\mathbf{s} T_A(\mathbf{s}) T_A(\mathbf{s} - \mathbf{b}) S(\mathbf{b}, \mathbf{s})}{\int d^2\mathbf{s} T_A(\mathbf{s}) T_A(\mathbf{s} - \mathbf{b})}$$

where  $S(\mathbf{b}, \mathbf{s})$  is the survival probability.


# $R_{AA}$ in the perturbative scenario



# $R_{AA}$ in the perturbative scenario




# $R_{AA}$ in the lattice scenario







# $R_{AA}$ in the lattice scenario



# $R_{AA}$ in the lattice scenario



# $R_{AA}$ in the lattice scenario



# Comments

- We expect our model to be more reliable at intermediate/large centralities since the condition  $E \gg \Gamma$  is more likely to be fulfilled.

# Comments

- We expect our model to be more reliable at intermediate/large centralities since the condition  $E \gg \Gamma$  is more likely to be fulfilled.
- The effect of the gap is around a factor of 2 in  $R_{AA}$  in all cases.

# Comments

- We expect our model to be more reliable at intermediate/large centralities since the condition  $E \gg \Gamma$  is more likely to be fulfilled.
- The effect of the gap is around a factor of 2 in  $R_{AA}$  in all cases.
- The effect is larger in the lattice scenario. The reason is that the gap computed from the formula  $2M_B - M_{\Upsilon(1S)}$  is larger than the gap computed from a Coulombic potential.

# Conclusions

- The energy gap between singlets and octets makes the medium evolution of bound states very different in QCD compared to QED.

# Conclusions

- The energy gap between singlets and octets makes the medium evolution of bound states very different in QCD compared to QED.
- In the limit  $E \gg \Gamma$  the evolution of the density matrix can be simplified to a rate equation.



# Conclusions

- The energy gap between singlets and octets makes the medium evolution of bound states very different in QCD compared to QED.
- In the limit  $E \gg \Gamma$  the evolution of the density matrix can be simplified to a rate equation.
- Using the  $E \gg \Gamma$  approximation and the large- $N_c$  limit, we have checked the impact of considering the gap in phenomenological applications.

# Conclusions

- The energy gap between singlets and octets makes the medium evolution of bound states very different in QCD compared to QED.
- In the limit  $E \gg \Gamma$  the evolution of the density matrix can be simplified to a rate equation.
- Using the  $E \gg \Gamma$  approximation and the large- $N_c$  limit, we have checked the impact of considering the gap in phenomenological applications.
- We see a substantial effect. Both in a HTL inspired scenario and in a Lattice QCD inspired scenario.