



Contribution ID: 246

Type: Poster

## Overlap integrals of di-neutron cluster state and shell model state

*Tuesday 7 September 2021 11:43 (1 minute)*

Di-neutron correlations are extensively explored in recent experiments, and the enhancement of the spatial localization of the pair of the neutrons ( $n$ ) has been confirmed at the nuclear surface in the light neutron-excess systems, such as  $^{11}\text{Li}$  and  $^{19}\text{B}$ . The spatial localization of two neutrons, which is called “the di-neutron correlations”, are also investigated theoretically by employing the three-body model with core +  $n + n$ . The analysis of the di-neutron correlation gives an important key to elucidate the surface structure of neutron stars. If the di-neutron configuration is stabilized in the nuclear medium, there is a possibility of the formation of new phase with the di-neutron condensation at the surface of the neutron star.

If we consider the three-body system of core plus two neutrons, the spatial localization of two neutrons corresponds to the formation of “di-neutron cluster” around the core nucleus. On the other hand, the valence neutrons usually perform the independent particle motion around the core nucleus, and the ground state of the normal nucleus is explained by the so-called the nuclear shell model. The shell model configuration and the di-neutron one seem to be very different structure intuitively but these two configurations are non-orthogonal, and there is a finite amplitude of the di-neutron cluster component even if the pure shell model state is realized. Thus, in order to understand the feature of the di-neutron cluster more deeply, it is important to evaluate the overlap integral of the di-neutron cluster state and the shell model state, which is a measure of the non-orthogonal amplitude of these two different states.

We have developed a new formula to evaluate the overlap integral of the cluster and shell-model configurations, and the formula is applied to the core +  $n + n$  systems. In this report, we will report the systematic feature of the overlap integrals of the di-neutron cluster state (core +  $2n$ ) and the shell model state (core +  $n + n$ ) with a variation of the core mass number. In particular, we will discuss the enhancements of the overlap integral in connection to the single particle orbits of the valence two neutrons in the shell model states.

**Author:** ITO, Makoto (Department of Pure and Applied Physics, Kansai University)

**Co-authors:** NAKAMOTO, riu (department of pure and applied physics, kansai university); Mr UEDA, Eiju (Department of Pure and Applied Physics, Kansai University); Prof. SHIMIZU, Noritaka (Center for Nuclear Study, The university of Tokyo)

**Presenter:** ITO, Makoto (Department of Pure and Applied Physics, Kansai University)

**Session Classification:** Poster Session I

**Track Classification:** Nuclear and particle astrophysics