



Contribution ID: 146

Type: **Talk**

Search for K^+ decays to a lepton and invisible particles at NA62 (12+3)

Wednesday 8 September 2021 15:55 (15 minutes)

The NA62 experiment at CERN reports searches for $K^+ \rightarrow e^+N$, $K^+ \rightarrow \mu^+N$ and $K^+ \rightarrow \mu^+vX$ decays, where N and X are massive invisible particles, using the 2016-2018 data set.

The N particle is assumed to be a heavy neutral lepton, and the results are expressed as upper limits of $O(10^{-9})$ and $O(10^{-8})$ of the neutrino mixing parameter $|U_{e4}|^2$ and $|U_{\mu 4}|^2$, improving on the earlier searches for heavy neutral lepton production and decays in the kinematically accessible mass range.

The X particle is considered a scalar or vector hidden sector mediator decaying to an invisible final state, and upper limits of the decay branching fraction for X masses in the range 10-370 MeV/ c^2 are reported for the first time, ranging from $O(10^{-5})$ to $O(10^{-7})$.

An improved upper limit of 1.0×10^{-6} is established at 90% CL on the $K^+ \rightarrow \mu^+vvv^-$ branching fraction.

Primary author: SCHUCHMANN, Simone (Johannes Gutenberg University Mainz)

Presenter: SCHUCHMANN, Simone (Johannes Gutenberg University Mainz)

Session Classification: Flavour physics - CKM and beyond

Track Classification: Flavour physics - CKM and beyond