

Search for CP-odd $t\bar{t}H$ production in the single-lepton $H \rightarrow b\bar{b}$ decay channel


Luis Coelho¹, Emanuel Gouveia², Ana Luísa Carvalho³,
Ricardo Gonçalo¹, António Onofre⁴

¹LIP, UC; ²LIP, UMinho; ³LIP, IST; ⁴LIP, UMinho, CF-UM-UP



Introduction

- ▶ The SM Higgs boson is a scalar with no CP-violating interactions.
- ▶ The CP properties of the top quark Yukawa interaction can be directly probed through $t\bar{t}H$ and tH processes.
- ▶ Top Yukawa interaction was parametrised as



$$\mathcal{L}_{t\bar{t}H} = -k'_t y_t \phi \bar{\psi} (\cos \alpha + i\gamma_5 \sin \alpha) \psi$$

- ▶ Pure CP-even (CP-odd) coupling correspond to $\alpha = 0^\circ$ (90°).

Event Selection

- ▶ Events divided according to jet and b-tag multiplicity.
- ▶ 70% b-tagging efficiency.
- ▶ Signal-depleted regions used to constrain backgrounds from data.

Region	Leptons	Jets	B-tag	Higgs candidates
$SR_{inc}^{\geq 6j \geq 4b}$	= 1	≥ 6	≥ 4	-
$SR_{inc}^{\geq 5j \geq 4b}$		= 5		-
$SR_{boosted}$		≥ 4	-	≥ 1


▶ Boosted Higgs Candidate:

- Large-R jet with $p_T > 300$ GeV and $m \in (100 - 140)$ GeV, containing two b-tagged jets.


Analysis Strategy

- ▶ One BDT is trained to separate signal from background.
- ▶ Inclusive signal regions are divided according to the output of the classification BDTs as control region or signal region depending on the signal-to-background ratio.

Background composition




- ▶ Another BDT is trained to distinguish CP-even from CP-odd $t\bar{t}H$ events.
- ▶ Lab-frame observables and angular variables calculated in specific frames can be used to discriminate between CP scenarios [1, 2].
- ▶ In the inclusive signal regions, the CP BDT is used as input to the fit. In the boosted channel, the classification BDT is used.



Measurement of the CP-Mixing Angle

- ▶ Analysis based on a binned likelihood $\mathcal{L}(\alpha, k'_t, \theta)$.
- ▶ Number of events in a given bin depends on α , on k'_t and on the set of nuisance parameters θ .
- ▶ Parameters are allowed to vary, the best-fit α and k'_t maximise the likelihood.
- ▶ 0.26π (47°) expected uncertainty on the signal.
- ▶ Good agreement with the background model.



Contributions

- ▶ tH signal parameterization as function of the mixing angle α and the coupling strength k'_t .
- ▶ Background-only fit to evaluate the efficacy of the background model.

References

- [1] D. Azevedo, A. Onofre, F. Filthaut, and R. Gonçalo. CP tests of Higgs couplings in $t\bar{t}H$ semileptonic events at the LHC. *Phys. Rev. D*, 98(3):033004, 2018.
- [2] A. Ferroglia, M. C. Fiolhais, E. Gouveia, and A. Onofre. Role of the $t\bar{t}H$ rest frame in direct top-quark Yukawa coupling measurements. *Phys. Rev. D*, 100(7):075034, 2019.