

Development of a next-generation detector concept to detect astrophysical gamma-rays

Responsables: Ruben Conceição and Bernardo Tomé;

- Matilde Tubal, FCUL at UL;
- Tatiana Mendes, FCUP at UP.



How to detect gamma rays?




Fig1: Satellites for high energy gamma-rays and Cherenkov telescopes or EAS arrays for very high energy gamma-rays;

How to measure the X_{\max}






Fig2: Relation between the X_{\max} and a constant “ c ”, which measures the shower front plane curvature;

$$X_{\max}^1 = B_1 + \gamma_1 c$$

Our goal


1. Calibrate with photon and proton;
2. Reconstruct with photon, proton and iron;
3. Check the results;
4. Conclusion;




Our results

Calibrations:

- Photon




- Proton




Our results

photon calibration




Photon

Xmax Bias vs Xmax:




Xmax Resolution vs Sem:




Sem - electromagnetic energy at the ground - proportional to the shower energy

Proton

Xmax Bias:




Xmax Resolution vs Sem:



Separation
between proton
and iron


Our results

proton calibration




Proton

Xmax Bias:




Xmax Resolution vs Sem:




Separation
between proton
and iron

Proton - Xmax reconstructed distributions


Sem=199.526



Sem=501.187



Sem=1995.26




Summary

- It is possible to correlate the X_{max} with the shower curvature (c) not only for photon induced showers but also for hadronic (proton and iron) showers
- The calibration with photon showers cannot be used for proton induced showers
 - A bias on X_{max} appears
- The calibration with proton showers removes the bias on X_{max} but the obtained resolution is too poor to distinguish between proton and iron showers
- The X_{max} reconstructed distributions have an healthy Gaussian shape suggesting that the method is working but the shower curvature reconstruction (c) should be revisited


Backup slides

Photon/Photon

Calibration plot:




Bias X_{\max} :




Photon/Photon


Xmax Bias vs Sem:




Xmax Resolution vs Sem:




Sem = 199.526



Sem = 501.187




Sem = 1995.26




Photon/Proton

Calibration plot:




Bias X_{\max} :





Photon/Proton


Bias Sem:




Resolution Sem, (0-200):




Sem = 199.526



Sem = 501.187




Sem = 1995.26




Photon/Iron

Calibration plot:




Bias Xmax:





Photon/Iron


Bias Sem:




Resolution Sem, (0-200):




Sem = 199.526



Sem = 501.187




Sem = 1995.26




Proton/Photon

Calibration plot:





Bias Xmax:




Proton/Photon

Bias Sem:




Resolution Sem:




Proton/Photon


Sem=199.526



Sem=501.187




Sem=1995.26




Proton/Proton

Calibration plot:





Bias Xmax:




Proton/Proton

Bias Sem:




Resolution Sem:




Proton/Proton


Sem=199.526



Sem=501.187




Sem=1995.26




Proton/Iron

Calibration plot:





Bias X_{\max} :




Proton/Iron

Bias Sem:




Resolution Sem:




Proton/Iron

Sem=199.526



Sem=501.187



Sem=1995.26

